

CONCENTRATION INEQUALITIES WITH
EXCHANGEABLE PAIRS

A DISSERTATION
SUBMITTED TO THE DEPARTMENT OF STATISTICS
AND THE COMMITTEE ON GRADUATE STUDIES
OF STANFORD UNIVERSITY
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

Sourav Chatterjee
February 2008

© Copyright by Sourav Chatterjee 2008
All Rights Reserved

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

Persi Diaconis Principal Adviser

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

Amir Dembo

I certify that I have read this dissertation and that, in my opinion, it is fully adequate in scope and quality as a dissertation for the degree of Doctor of Philosophy.

David O. Siegmund

Approved for the University Committee on Graduate Studies.

Abstract

The purpose of this dissertation is to introduce a version of Stein’s method of exchangeable pairs to solve problems in measure concentration. We specifically target systems of dependent random variables, since that is where the power of Stein’s method is fully realized. Because the theory is quite abstract, we have tried to put in as many examples as possible. Some of the highlighted applications are as follows: (a) We shall find an easily verifiable condition under which a popular heuristic technique originating from physics, known as the “mean field equations” method, is valid. No such condition is currently known. (b) We shall present a way of using couplings to derive concentration inequalities. Although couplings are routinely used for proving decay of correlations, no method for using couplings to derive concentration bounds is available in the literature. This will be used to obtain (c) concentration inequalities with explicit constants under Dobrushin’s condition of weak dependence. (d) We shall give a method for obtaining concentration of Haar measures using convergence rates of related random walks on groups. Using this technique and one of the numerous available results about rates of convergence of random walks, we will then prove (e) a quantitative version of Voiculescu’s celebrated connection between random matrix theory and free probability.

Acknowledgements

First of all, I would like to thank my parents for bringing me up the way they did, because, despite all the deficiencies in my character I am sure it is immensely better than what I could have managed on my own. They don't know any math at all, but they have the sincere respect for academics that is so pleasantly ubiquitous in the land from where I hail. It is also worthy of note, dear reader, that they love me more than anything else in the universe.

Having said that, let me add that my character and enthusiasm alone would not have made me a mathematician. There is one person who is overwhelmingly responsible for that, and that is my adviser, Persi Diaconis. I recall that Persi once told me, towards the beginning of my Ph.D. quest, that his job was to take me to the “top”. Well, there is still a long way to go, but I am grateful to him for at least showing me with his own attitude what the “top” demands and bringing it closer to the earth. I also thank him for numerous other things, not the least for reading this thesis with an unbelievable amount of diligence and care.

I would also like to thank professors Amir Dembo and David Siegmund for agreeing to be readers for this dissertation, and also for the numerous things I learned over the past three years from these two astute mathematicians. Amir has been particularly kind to me on many occasions, sparing a lot of his time on academic discussions and giving advice on various issues.

Charles Stein has been an inspiration in all the work that I did during my Ph.D.

Besides that, I thank all the other professors and staff in this wonderful department who have never flinched about answering a question from a nagging graduate student — something that is embedded in the culture of Sequoia Hall.

I am grateful to professor Arup Bose from the Indian Statistical Institute, Kolkata, for initiating my research career by helping me write my first real paper.

It is imperative that I thank all the professors who answered my various questions by e-mail; in that respect, I am particularly grateful to professor Michel Ledoux. The other members of this illustrious list include Michel Talagrand, Boguslaw Zegarlinski, Kenneth Alexander and David Aldous. I hope I have not missed anyone out!

Thanks are also due to Dadabhai (Rajib) and Boudi (Mukta), and of course, little Roompi, for taking care of me in various ways in the last three years. They add a touch of sanity to my world that is otherwise mired in academics.

Coming to friends, I have a few of them to whom I owe a bit (not money, fortunately). First of all, there is Soumik Pal, my closest confidante and friend, from whom I learned about life as much as I did about mathematics. He never fails to amaze me with his mathematical originality at regular intervals of time; for instance, I should mention that he has a beautiful alternative martingale argument for one of the basic theorems in this dissertation (Theorem 3.3).

Then there is Pinaki Biswas, and he may or may not be surprised to see his name in this list; but I am grateful to him, if for nothing else, then for a friendship that has lasted seventeen years now, and that is something too valuable by itself to discount.

I must also thank my three closest friends at Stanford — George Chang, Nancy Ruonan Zhang, and Brit Turnbull. They are the ones most responsible for making my life livable in this desolate rural countryside.

Finally, there is one person from the past, an old retired clerk to whom I owe more than what I realized when he was alive. He was my childhood friend; in the long hours that we used to spend together, he introduced me to everything ranging from Euclid to atheism, from simultaneous equations to Shakespeare; in short, everything that inspired me to think and to dream. I would like to dedicate this thesis to the memory of my grandfather, the late Shri Tarapada Chatterjee.

Contents

Abstract	iv
Acknowledgements	v
1 Introduction	1
1.1 Summary of thesis	2
2 Review of existing literature	7
2.1 Hoeffding type inequalities	7
2.2 Talagrand's concentration inequalities	11
2.3 Logarithmic Sobolev inequalities	13
2.4 Other entropy based methods	17
2.5 Geometric measure concentration	21
2.6 Concentration on groups	23
2.7 Stein's method of exchangeable pairs	26
3 Theory and examples: Part I	28
3.1 A variance formula	29
3.2 An exponential inequality	31
3.3 Example: Curie-Weiss model	33
3.4 Validity of naive mean field equations	36
3.5 Example: Least squares for the Ising model	42
3.6 An inequality for self-bounded functions	49
3.7 Example: Matching problem and Spearman's footrule	52

3.8	Inequalities for unbounded differences	55
3.9	A refinement of the exponential inequality	60
3.10	Application to spin glasses	62
4	Theory and examples: Part II	69
4.1	Explicit construction of $F(X, X')$	70
4.2	Concentration under weak dependence	74
4.3	Example: Spin systems	79
4.4	Example: Graph colorings	82
4.5	Concentration of Haar measures	84
4.6	Application to random matrices and freeprobability	90
	Bibliography	95

Chapter 1

Introduction

The theory of concentration inequalities tries to answer the following question: Given a random variable X taking value in some measure space $\mathcal{X}$ (which is usually some high dimensional Euclidean space), and a measurable map $f : \mathcal{X} \rightarrow \mathbb{R}$, what is a good explicit bound on $\mathbb{P}\{|f(X) - \mathbb{E}f(X)| \geq x\}$? Exact evaluation or accurate approximation is, of course, the central purpose of probability theory itself. In situations where this is not possible, concentration inequalities aim to do the next best job.

A bound is *good* if it is sufficiently rapidly decreasing; gaussian bounds are usually considered satisfactory. The reasons for insisting on good bounds (as opposed to Chebychev type bounds) are theoretical as much as practical. In fact, the theoretical interest often supercedes the practical aspect, because in spite of all the activity, concentration bounds often give bad numbers when calculated numerically. Theoretically their importance stems, in large part, from the Bonferroni inequality: If we know that if a collection of events $\{A_k\}_{1 \leq k \leq n}$ are so rare that $\mathbb{P}\{A_k\} \leq e^{-cn}$ for each k for some fixed constant c , then $\mathbb{P}\{\cup_k A_k\} \leq ne^{-cn}$, which is again “small”, since the e^{-cn} term “kills” the n . Contrary to what someone unfamiliar with the literature might feel, this seemingly crude technique has been successful in establishing surprisingly efficient results, mainly because “rare events are often approximately disjoint”.

This was at the center of the earliest line of thought about controlling the suprema of empirical processes, developed mainly by David Pollard and others in the eighties (see Pollard [87] for an exposition). Subsequent developments in the nineties,

based on Talagrand’s concentration inequalities [107, 109, 110] and the more recent “entropy method” of Ledoux [61] and Massart [69], are more subtle. They will be discussed later in detail. All in all, concentration inequalities form the backbone of this very important branch of modern theoretical statistics and machine learning. In return, empirical process theory forms the most prominent area of application for concentration bounds.

Concentration inequalities are also used in theoretical computer science, random matrix theory and a variety of other fields, for reasons more or less the same as mentioned before. Often, applications get obscured because they were not mentioned in the abstract or the keywords, and that is a sign that concentration inequalities are rapidly attaining the status of standard tools like the Borel-Cantelli lemmas in the classical probability literature. There may soon be a time when researchers will start using Talagrand’s inequalities without explicit reference in the abstract.

The theory of concentration inequalities for functions of independent random variables, some of which was described above, has reached a high level of sophistication by now. However, concentration inequalities for functions of dependent random variables are still hard to get, the main tools being logarithmic Sobolev and transportation cost inequalities. One shortcoming of these methods is that explicit constants are very hard or almost impossible to get. We shall discuss these methods in detail in the next chapter.

The main purpose of this thesis is to construct a modification of Stein’s method of exchangeable pairs, which is a well-known tool from probability theory, to derive concentration inequalities with explicit constants for functions of dependent random variables. We postpone a discussion of Stein’s method until Section 2.7.

1.1 Summary of thesis

We now give a brief chapter by chapter description of this thesis. The theory is too abstract and requires too much notation to describe in this brief introductory discussion. Instead, we shall present some theorems which were obtained as applications of the theory, for the purpose of enticing the prospective reader to delve deeper.

In Chapter 2, we shall review the major existing results from the concentration inequality literature, including the early martingale techniques, Talagrand's inequalities, and modern entropy based methods. We shall also describe the basic philosophy of Stein's method in the last section of this chapter.

Chapter 3 is the first part of our theory, which is the more “abstract” part. We shall develop our basic results in this chapter, and apply them to work out some simple examples involving dependent variables: In section 3.3, we shall obtain a simple and explicit concentration bound for $m - \tanh(\beta m)$, where m is the magnetization in the Curie-Weiss model of ferromagnetic interaction. By a generalization of this example, we shall derive in section 3.4 a broad condition under which the naive mean field equations — one of the standard tools from physics that is frowned upon by mathematicians (for good reasons) — is valid. In section 3.5, we shall work out an example about estimation of parameters in statistical models of dependent data. In section 3.7, we shall apply our techniques to obtain tail bounds of the correct order for (a) the number of fixed points in a random permutation, and (b) the Spearman's footrule distance between a random permutation and the identity. Finally, in section 3.10, we shall obtain a concentration result about the Sherrington-Kirkpatrick model of spin glasses which holds at all temperatures.

In Chapter 4, we shall develop some advanced tools (Lemma 4.1, Lemma 4.2) so that the theorems from Chapter 3 can be easily applied to more complex problems than the ones worked out in Chapter 3. Indeed, Lemma 4.2, in conjunction with Theorem 3.3, is probably the first general tool which allows one to use couplings to prove concentration inequalities. Couplings are widely used to establish rates of temporal and spatial decay of correlations; so it seems natural that there should be some result which makes them useful for concentration inequalities, too.

An application of these tools gives Theorem 4.3, which allows us to derive concentration inequalities for arbitrary functions in complicated models of dependent random variables, such as Ising type spin models and random proper colorings of finite graphs, under a Dobrushin type condition of weak dependence. We shall now state this theorem, after introducing some required notation.

Let Ω be a Polish space and let $f : \Omega^n \rightarrow \mathbb{R}$ be a function satisfying a Lipschitz

condition with respect to a generalized Hamming distance on Ω^n : For all $x, y \in \Omega^n$,

$$|f(x) - f(y)| \leq \sum_{i=1}^n c_i \mathbb{I}\{x_i \neq y_i\} \quad (1.1)$$

where $c_1, \dots, c_n$ are some fixed nonnegative constants. This just means that the value of the function does not change by more than c_i if the i^{th} coordinate is altered.

Let $X = (X_1, \dots, X_n)$ be an Ω^n -valued random variable with law μ . For any $x \in \Omega^n$, let $\bar{x}^i$ denote the element of Ω^{n-1} obtained by omitting the i^{th} coordinate of x . For each $i \leq n$ and $x \in \Omega^n$, let $\mu_i(\cdot | \bar{x}^i)$ denote the law of X_i given $\bar{X}^i = \bar{x}^i$. Finally, let us recall that for a square matrix A , the L^2 operator norm of A is defined as:

$$\|A\|_2 := \max_{\|y\|=1} \|Ay\|.$$

Then we have the following result from section 4.2:

Theorem 4.3 *Suppose $A = (a_{ij})$ is an $n \times n$ matrix with nonnegative entries and zeros on the diagonal such that for any i , and any $x, y \in \Omega^n$,*

$$d_{TV}(\mu_i(\cdot | \bar{x}^i), \mu_i(\cdot | \bar{y}^i)) \leq \sum_{j=1}^n a_{ij} \mathbb{I}\{x_j \neq y_j\},$$

where d_{TV} is the total variation distance on the space of probability measures on Ω . Suppose f satisfies the generalized Lipschitz condition (1.1). If $\|A\|_2 < 1$, we have

$$\mathbb{P}\{|f(X) - \mathbb{E}f(X)| \geq t\} \leq 2e^{-(1-\|A\|_2)t^2 / \sum_i c_i^2}$$

for each $t \geq 0$.

This is possibly the first result which gives a direct connection between the Dobrushin condition [35] from statistical physics and concentration inequalities. The log-Sobolev inequalities of Stroock & Zegarlinski [103, 104, 105] do not give explicit constants, and it is also not clear whether they extend to systems beyond the lattice.

Some applications of the above theorem, to spin systems and graph colorings, will be worked out in sections 4.3 and 4.4. In particular, we shall work out explicit

concentration bounds for the magnetization in the Ising model on an arbitrary graph at high temperature.

In section 4.5 of the same chapter, we shall derive a tool (Theorem 4.6) for obtaining the concentration of measures which are invariant under group actions. This is possibly the first result which gives an explicit connection between rates of convergence to stationarity for random walks on groups and the concentration of Haar measures. Using this result, we shall obtain the following quantitative bound related to random matrices and free probability theory, which again, is probably the first result of its kind. The limiting version of this result is quite well-known; following from a celebrated result of Voiculescu [114], it says, roughly, that the distribution of the eigenvalues of $M + N$, where M and N are two high dimensional hermitian matrices of the same order, is approximately determined by the eigenvalue distributions of M and N .

In the following theorem, the term “empirical distribution function of H ” is a commonly used random matrix jargon, which just means the probability distribution function on $\mathbb{R}$ which puts mass $1/n$ on each eigenvalue of the matrix H .

Theorem 4.10 *Let Δ_1 and Δ_2 be two $n \times n$ real diagonal matrices. Let U and V be independent Haar distributed random elements of $\mathcal{U}_n$, the group of all unitary matrices of order n . Let*

$$H = U\Delta_1U^* + V\Delta_2V^*,$$

and let F_H be the empirical distribution function of H . Then, for every $x \in \mathbb{R}$, $\text{Var}(F_H(x)) \leq \kappa n^{-1} \log n$. where κ is a universal constant not depending on n , Δ_1 , Δ_2 or x . Moreover we also have the concentration inequality

$$\mathbb{P}\{|F_H(x) - \mathbb{E}(F_H(x))| \geq t\} \leq 2 \exp\left(-\frac{nt^2}{2\kappa \log n}\right)$$

for every $t \geq 0$, where κ is the same as in the variance bound.

We think that it will be very hard to derive such a result using available techniques, because there is no independence, and gaussianity is involved in a complicated way, and the standard concentration results for gaussian measures are all with respect to

the Euclidean metric, which cannot give a universal bound like the above (bounds, if any, will involve Δ_1 and Δ_2 , and will be inefficient, because the function F_H is badly nonsmooth). On the other hand, our bound follows quite easily from the theory developed in this dissertation. We do not know, however, if it is of the correct order (even after discounting the $\log n$ factor).

Finally, let us mention some of the deficiencies of this dissertation. One shortcoming is in the range of examples. Although we have tried our best to provide as many as we can, it is probably not enough to cover all fields of interest; in particular, we have no examples from empirical process theory. Moreover, the author is not completely happy with the quality of some of the applications. For instance, the result about spin glasses seems to have a wide scope, but the author has been frustrated by attempts to exploit it further. Another unfinished aspect is that we could not find good examples for the “unbounded differences” theorems of section 3.8.

It is also unfortunate that the technique in Chapter 4.2 has only been applied to get Theorems 4.3 and 4.6. We feel that the scope of Lemma 4.2, which allows us to get concentration bounds using couplings, extends much beyond that.

Finally, a major incompleteness, which is a weakness of the idea of concentration itself, is the lack of lower bounds.

We shall attempt to overcome these and other deficiencies in future work.

Chapter 2

Review of existing literature

This chapter will be devoted to the discussion of the main existing tools for proving concentration bounds. We apologize in advance for unjust omissions, if any. Most of the material in this chapter, except for the very recent developments, is taken from the wonderful monograph by Michel Ledoux [61]. In fact, in many places we have kept the notation and even the language, intact.

While we shall attempt to give a comprehensive summary of the main theoretical results, we shall, in general, refrain from discussing applications in this chapter. The main reason being, applications invariably entail some lengthy background, the discussion of which would be a digression from the central theme of this chapter. This is the same reason why we shall not discuss concentration inequalities in empirical process theory. Of course, generous references will be provided.

In the last section of this chapter, we shall discuss the basics of Stein's method.

2.1 Hoeffding type inequalities

The Azuma-Hoeffding martingale inequality [52, 7] remained the last word in concentration for a very long time, culminating in the bounded difference inequality, observed by Schechtman [97] and also by Shamir & Spencer [99], and its extensive popularity among combinatorialists and discrete mathematicians following the expository work of McDiarmid [71].

In its most widely used form, the Azuma-Hoeffding inequality goes as follows:

Theorem 2.1 [Azuma-Hoeffding inequality [52, 7]] *Let $\{X_i\}_{1 \leq i \leq n}$ be a martingale difference sequence adapted to some filtration. Suppose $c_1, \dots, c_n$ are constants such that $|X_i| \leq c_i$ almost surely for each i . Then for all $t \geq 0$ we have*

$$\mathbb{P}\left\{\max_{1 \leq k \leq n} \sum_{i=1}^k X_i \geq t\right\} \leq \exp\left(-\frac{t^2}{2 \sum_{i=1}^n c_i^2}\right).$$

Before stating the bounded difference inequality, we must state the most “accurate” version of the Hoeffding inequality for independent summands, which was established by Bennett [11].

Theorem 2.2 [Bennett’s inequality [11]] *Let $Y_1, \dots, Y_n$ be independent real-valued random variables bounded in magnitude by some constant C . Let $S = \sum_{i=1}^n Y_i$. Then, for every $t \geq 0$,*

$$\mathbb{P}\{S \geq \mathbb{E}(S) + t\} \leq \exp\left\{-\frac{\sigma^2}{C^2} h\left(\frac{Ct}{\sigma^2}\right)\right\},$$

where $h(u) = (1 + u) \log(1 + u) - u$, and $\sigma^2 = \sum_{i=1}^n \mathbb{E}(Y_i^2)$.

Originally proved by Bernstein for Bernoulli random variables, this is also sometimes called Bernstein’s inequality.

The bounded difference inequality, stated in its simplest form, is the following powerful corollary of the Azuma-Hoeffding inequality:

Theorem 2.3 [Bounded difference inequality [97, 99, 71]] *Let $Z = f(X_1, \dots, X_n)$ be a function of independent random variables $X_1, \dots, X_n$. Let X'_i be an independent copy of X_i , $i = 1, \dots, n$. Suppose $c_1, \dots, c_n$ are constants such that for each i ,*

$$|f(X_1, \dots, X_{i-1}, X'_i, X_{i+1}, \dots, X_n) - f(X_1, \dots, X_n)| \leq c_i \quad a.s.$$

Then, for any $t \geq 0$ we have

$$\mathbb{P}\{Z - \mathbb{E}(Z) \geq t\} \leq \exp\left(-\frac{2t^2}{\sum_{i=1}^n c_i^2}\right).$$

The above inequality, though widely useful, does not in general convey the theoretical essence of the situation, because the bounds on the differences do not reflect the typical size of the differences in many interesting problems. A better result, from the conceptual point of view, is the following inequality, first discovered by Efron and Stein [40], which has since come to be known as the Efron-Stein inequality. The present version is due to Steele [100]:

Theorem 2.4 [Efron-Stein inequality [40, 100]] *Keeping the notation exactly as in the previous theorem, we have*

$$\text{Var}(Z) \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E}[(f(X_1, \dots, X_{i-1}, X'_i, X_{i+1}, \dots, X_n) - f(X_1, \dots, X_n))^2].$$

The Efron-Stein inequality has been used to bound variances in many complicated problems. For a very recent application, one can see the work of Reitzner [90] on random polytopes. However, the inequality is not guaranteed to give a bound of the correct order. For instance, the actual variance of the number of k -gons in an Erdős-Rényi random graph is less than the Efron-Stein bound by a factor of k (Cf. [19], section 6). Also, it is only a variance bound and gives no useful information about the tail behavior.

All of the above results are based on martingale methods. The limitations of martingale techniques were gradually recognized in the late eighties, and people started looking for alternatives. In recent years, however, there has been some kind of a minor resurgence of interest in extending the old martingale arguments. One of the more successful efforts has been the so-called “divide and conquer martingale” method of Kim & Vu [58]. We refer to this paper for further references to the current literature around the martingale method.

A significant discovery was recently made by Boucheron, Lugosi & Massart [19], who established an exponential version of the Efron-Stein inequality. In our opinion, this work has resulted in the completion of the quest for maximum efficiency in concentration via Hoeffding type inequalities.

The methodology used to prove the exponential Efron-Stein is based on the “entropy method” introduced by Ledoux [61] and Massart [69], and generalized by

Boucheron, Lugosi & Massart in [18]. This powerful new method is based on the modified log Sobolev approach, to be discussed in section 2.4.

Before stating the exponential Efron-Stein inequality, we have to introduce some notation.

Let $X_1, \dots, X_n$ be independent random variables taking values in a measurable space $\mathcal{X}$. Denote by X_1^n the vector of these n random variables. Let $f : \mathcal{X}^n \rightarrow \mathbb{R}$ be a measurable map. Let $Z = f(X_1, \dots, X_n)$. Let $X'_1, \dots, X'_n$ denote independent copies of $X_1, \dots, X_n$, and let $Z^{(i)} = f(X_1, \dots, X_{i-1}, X'_i, X_{i+1}, \dots, X_n)$.

Define the random variables V_+ and V_- by

$$V_+ = \mathbb{E} \left[\sum_{i=1}^n (Z - Z^{(i)})^2 \mathbb{I}_{Z > Z^{(i)}} \middle| X_1^n \right]$$

and

$$V_- = \mathbb{E} \left[\sum_{i=1}^n (Z - Z^{(i)})^2 \mathbb{I}_{Z < Z^{(i)}} \middle| X_1^n \right].$$

The following result appears as Theorem 2 in [19]:

Theorem 2.5 [Exponential Efron-Stein inequality] *For all $\theta > 0$ and $\lambda \in (0, 1/\theta)$,*

$$\log \mathbb{E} [\exp(\lambda(Z - \mathbb{E}(Z)))] \leq \frac{\lambda\theta}{1 - \lambda\theta} \log \mathbb{E} \left[\exp \left(\frac{\lambda V_+}{\theta} \right) \right].$$

On the other hand, we have for all $\theta > 0$ and $\lambda \in (0, 1/\theta)$,

$$\log \mathbb{E} [\exp(-\lambda(Z - \mathbb{E}(Z)))] \leq \frac{\lambda\theta}{1 - \lambda\theta} \log \mathbb{E} \left[\exp \left(\frac{\lambda V_-}{\theta} \right) \right].$$

The paper [19] has several nice applications of this result, including applications to empirical processes and subgraph counts. In particular, the authors prove that Talagrand's famous convex distance inequality (to be discussed in the next section) is a corollary of the exponential Efron-Stein inequality.

The more recent paper [20] on generalized moment inequalities in the same spirit as the exponential Efron-Stein inequality, is also worthy of note.

2.2 Talagrand's concentration inequalities

This section is devoted to the discussion of the deep investigation of Michel Talagrand, which climaxed in a series of intricate and powerful results about concentration of measure in product spaces [107, 109, 110]. These have since come to be known as Talagrand's concentration inequalities. Rooted in abstract geometric formulation, Talagrand's techniques have found wide applications in fields ranging from combinatorial optimization (e.g. traveling salesman problem in [107]) to random matrix theory [50, 2].

Before we begin our discussion, let us introduce some basic notation: Throughout, we consider a product probability measure μ^n on a product space $\mathcal{X}^n$. For any vector $c = (c_1, \dots, c_n) \in \mathbb{R}_+^n$, the generalized Hamming metric d_c is defined on $\mathcal{X}^n$ as

$$d_c(x, y) := \sum_{i=1}^n c_i \mathbb{I}\{x_i \neq y_i\}. \quad (2.1)$$

Thus, the bounded difference inequality says that if $f : \mathcal{X}^n \rightarrow \mathbb{R}$ is a function such that $|f(x) - f(y)| \leq d_c(x, y)$ for all x and y , then $\mu^n\{f - \int f d\mu^n \geq t\} \leq \exp(-t^2/2\|c\|^2)$, where $\|c\|^2 := \sum_{i=1}^n c_i^2$.

A fundamental weakness of this inequality and its early variants is the following: They only allow us to consider functions f which satisfy the Lipschitz condition for some fixed c . It was soon realized that a lot of open questions in measure concentration were about functions which do not satisfy (2.1), but rather, obey

$$f(x) - f(y) \leq \sum_{i=1}^n c_i(x) \mathbb{I}\{x_i \neq y_i\} \quad (2.2)$$

or some variant of this, for some vector field $c : \mathcal{X}^n \rightarrow \mathbb{R}^n$ with uniformly bounded norm. (The asymmetry in the above expression is often a help rather than hindrance in applications.) It is thus desirable to have a Hoeffding type bound for such functions based on $\sup\{\|c(x)\|^2 : x \in \mathcal{X}^n\}$. Talagrand's famed convex distance inequality is, in essence, a generalization of this idea in the abstract geometric setting.

Given a set $A \subseteq \mathcal{X}^n$, let us define the “convex distance” of a point x to A as

$$D_A(x) := \sup_{\|c\|=1} d_c(x, A),$$

where $d_c(x, A) = \inf_{y \in A} d_c(x, y)$. Note that $D_A(x) = d_{c(x)}(x, A)$ for some $c(x)$ depending on x if the supremum is attained (which is usually the case).

The following version of Talagrand’s main result is taken from Ledoux [61], Theorem 4.6:

Theorem 2.6 [Talagrand’s convex distance inequality [107]] *For every measurable non-empty subset A of $\mathcal{X}^n$, and for every product probability measure μ^n on $\mathcal{X}^n$,*

$$\int e^{D_A(x)^2/4} \mu^n(dx) \leq \frac{1}{\mu^n(A)}.$$

In particular, for every $t \geq 0$,

$$\mu^n\{D_A \geq t\} \leq \frac{1}{\mu^n(A)} e^{-t^2/4}.$$

To connect this with concentration problems for specific functions, the usual route is the following: Given a function f with median m_f (that is, $\mu^n\{f \geq m_f\} \geq 1/2$ and $\mu^n\{f \leq m_f\} \geq 1/2$), let $A = \{x : f(x) \leq m_f\}$. Then $\mu^n(A) \geq 1/2$. Next, find a function $r(t)$ such that $f(x) - m_f > t$ implies $D_A(x) > r(t)$. It is then easy to see that

$$\mu^n\{f - m_f > t\} \leq \mu^n\{D_A > r(t)\} \leq 2e^{-r(t)^2/4}.$$

The abstract formulation allows us to go beyond Lipschitz functions, in the following way (which is also the usual method of applying the convex distance inequality): Given a function f with median m_f , we have to find a function $r(t)$ having the following property: Whenever $f(x) \geq m_f + t$ and $f(y) \leq m_f$, there exists some $c = c(x) \in \mathbb{R}^n$ such that $\|c\| = 1$ and $\sum_{i=1}^n c_i(x) \mathbb{I}\{x_i \neq y_i\} > r(t)$. In the particular case when $r(t) = t$, we get the condition $f(x) - f(y) \leq \sum_{i=1}^n c_i(x) \mathbb{I}\{x_i \neq y_i\}$, which was stated as the initial motivation for this discussion. However, we may have $r(t) \neq t$ also, as demonstrated in Talagrand’s bound for the concentration of the longest

increasing subsequence in a random permutation ([107], section 7.1).

As is usual in geometric measure concentration (to be discussed in section 2.5), Talagrand's inequality gives concentration around the median rather than the mean, but that is a minor inconvenience.

A very important and striking consequence of Theorem 2.6 is the following:

Corollary 2.7 [Talagrand [107]] *For every product probability μ^n on $[0, 1]^n$, every convex 1-Lipschitz function f on $\mathbb{R}^n$, and every $t \geq 0$,*

$$\mu^n\{|f - m_f| \geq t\} \leq 4e^{-t^2/4},$$

where m_f is the median of f for μ^n .

The applications of Talagrand's convex distance inequality are too diverse to summarize in a few paragraphs. Many of the most striking examples were worked out by Talagrand himself in his landmark paper [107], including applications to the traveling salesman problem and first passage percolation. Applications to the concentration of spectral measures of random matrices were made by Guionnet & Zeitouni [50]. Concentration of individual eigenvalues of gaussian random matrices was established using Theorem 2.6 in a short but remarkable paper by Alon, Krivelevich & Vu [2].

Another important result of Talagrand is the “control by several points” method, which we shall not discuss here. For a description of this technique, one can look at Talagrand's original paper [107] or section 4.3 in Ledoux' book [61]. Talagrand used this method to prove his celebrated Bernstein type bound for empirical processes. This result is presented in a nice way as Theorem 7.4 in [61].

2.3 Logarithmic Sobolev inequalities

Logarithmic Sobolev inequalities were introduced by Gross [48] as the infinitesimal version of hypercontractivity in quantum field theory. They soon became a tool of fundamental importance, and the last three decades have seen vigorous activity surrounding the theory and applications of log-Sobolev inequalities. It is not our purpose

to go deeply into all that; we shall be only concerned with their relevance in measure concentration. For other applications in probability and statistical mechanics, one can look at the lecture notes by Guionnet and Zegarlinski [49] which concentrate on physical applications, and the paper by Diaconis and Saloff-Coste [32] on applications of log-Sobolev inequalities to finite Markov chains. An excellent survey of known mathematical results can be found in [4].

Suppose $X_0, X_1, \dots$, is a stationary reversible Markov chain on some space $\mathcal{X}$. The *Dirichlet form* corresponding to this chain (or more appropriately, corresponding to the associated kernel) is a functional $\mathcal{E}$ defined on the space of all pairs f, g of maps from $\mathcal{X}$ into $\mathbb{R}$ which satisfy $\mathbb{E}(f(X_0)^2) < \infty$ and $\mathbb{E}(g(X_0)^2) < \infty$. It is defined as

$$\mathcal{E}(f, g) := \frac{1}{2} \mathbb{E}[(f(X_1) - f(X_0))(g(X_1) - g(X_0))].$$

The kernel is said to satisfy a *logarithmic Sobolev inequality* with constant c if for all f such that $\mathbb{E}(f(X_0)^2) < \infty$, we have

$$\mathbb{E}\left(f(X_0)^2 \log \frac{f(X_0)^2}{\mathbb{E}f(X_0)^2}\right) \leq 2c\mathcal{E}(f, f).$$

Often, probability measures have natural reversible kernels associated with them. In such cases, instead of saying that the kernel satisfies a log-Sobolev inequality, we say that the probability distribution satisfies such an inequality. For instance, a distribution on $\mathbb{R}^n$ which has density $\rho(x)$ with respect to Lebesgue measure, is, under appropriate conditions, the stationary distribution of the Langevin diffusion process $(X_t)_{t \geq 0}$ with constant volatility matrix $\sqrt{2}I$ and drift $\nabla \log \rho(x)$. Instead of X_0 and X_1 we now take X_0 and X_h , where $h > 0$, but divide the right hand side by h in the definition of the log-Sobolev inequality. For the Langevin diffusion under suitable conditions, if we take $h \downarrow 0$, then $h^{-1}\mathcal{E}(f, f) \rightarrow \mathbb{E}(\|\nabla f(X_0)\|^2)$. This observation motivates the following definition: A probability measure μ on $\mathbb{R}^n$ is said to satisfy a logarithmic Sobolev inequality with constant c if for all locally Lipschitz f ,

$$\int f(x)^2 \log\left(\frac{f(x)^2}{\int f(u)^2 d\mu(u)}\right) d\mu(x) \leq 2c \int \|\nabla f(x)\|^2 d\mu(x).$$

To most people, this is the usual definition of a log-Sobolev inequality, but the earlier form is useful for discrete problems. The left hand side is usually called the entropy of f^2 with respect to μ , and is denoted by $\text{Ent}_\mu(f^2)$.

The single most important property of log-Sobolev inequalities is perhaps the tensorizing property: If $\mu_1, \dots, \mu_n$ are probability measures on $\mathbb{R}$ satisfying log-Sobolev inequalities with constants $c_1, \dots, c_n$, then the product measure $\mu_1 \otimes \dots \otimes \mu_n$ on $\mathbb{R}^n$ satisfies log-Sobolev inequality with constant $\max_i c_i$.

The connection between logarithmic Sobolev inequalities and concentration was made in an unpublished but now famous argument of I. Herbst. The following theorem, which summarizes the end result, is taken from Ledoux [61], Theorem 5.3:

Theorem 2.8 [Herbst's lemma] *Let μ be a probability measure on $\mathbb{R}^n$ satisfying a log-Sobolev inequality with constant c . Then, every 1-Lipschitz function $f : \mathcal{X} \rightarrow \mathbb{R}$ is integrable and for every $t \geq 0$, $\mu\{f \geq \int f d\mu + t\} \leq e^{-t^2/2c}$.*

Ledoux also presents a discrete version of the above result; the following occurs as Theorem 5.17 in [61]:

Theorem 2.9 *Let μ be the stationary distribution of a reversible Markov chain $\{X_k\}_{k \geq 0}$ on a countable set $\mathcal{X}$, which satisfies a logarithmic Sobolev inequality with constant c . Then, any $f : \mathcal{X} \rightarrow \mathbb{R}$ which satisfies*

$$\frac{1}{2} \sup_{x \in \mathcal{X}} \mathbb{E}((f(X_1) - f(X_0))^2 | X_0 = x) \leq 1,$$

also satisfies $\mu\{f \geq \int f d\mu + t\} \leq e^{-t^2/4c}$ for any $t \geq 0$.

One class of measures for which explicit log-Sobolev constants (and hence, concentration inequalities) are easily available are high dimensional probability measures with strictly log-concave density with respect to Lebesgue measure; this is a consequence of an important work of Bakry and Émery [9].

Theorem 2.10 [Bakry-Émery criterion] *Suppose μ is a measure on $\mathbb{R}^n$ with density $e^{-U(x)}$ with respect to Lebesgue measure, where $\text{Hess } U(x) \geq aI$ for some fixed constant $a > 0$, for all $x \in \mathbb{R}^n$. Then μ satisfies a log-Sobolev inequality with constant $1/a$.*

The result can be supplemented by the observation that if μ satisfies a logarithmic Sobolev inequality with constant c , then the measure defined by $d\nu = Z^{-1}e^V d\mu$ (where Z is the normalizing constant) satisfies a log-Sobolev inequality with constant $ce^{4\|V\|_\infty}$. This is the famous perturbation argument of Holley and Stroock [53]. However, this is not a very useful result in high dimensions, because $\|V\|_\infty$ blows up as the dimension increases.

It is to be noted that we do not necessarily have to go through the Herbst argument and Bakry-Émery's result to get the concentration inequality for strictly log-concave measures. Ledoux [61], pp. 39–41, has a direct proof.

However, in other cases, like spin systems on a lattice, there is no direct argument for concentration inequalities, and the route through the Herbst lemma must be used. In their important work [103, 104, 105], Stroock and Zegarlinski established that the Glauber dynamics associated with Ising and other spin models satisfy log-Sobolev inequalities at high temperature, wherever Dobrushin's condition of weak dependence holds. (See Georgii [45], Chapter 8, for the definition and examples for Dobrushin's condition. For a readable account of the work of Stroock and Zegarlinski, see the lecture notes [49].)

This implies, by Herbst's argument, that those spin systems satisfy concentration properties at high temperature. However, since explicit log-Sobolev constants are not known, it is not possible to get explicit concentration bounds along those lines.

This brings us to the point where we can state one of our achievements in this dissertation, which we already stated once in the Introduction: In section 4.2, we shall prove a concentration inequality with explicit constants for systems which satisfy Dobrushin's condition. We shall apply our result to get explicit concentration bounds for generalized Lipschitz functions of spins in Ising type models and uniformly chosen proper k -colorings of graphs with maximum degree $< k/2$.

For further information and references about the use of logarithmic Sobolev inequalities in the field of measure concentration, the reader is encouraged to look at the comprehensive survey [60]. For further details about their applications to spin systems, as well as some easy proofs of known results, one can look at [62].

2.4 Other entropy based methods

Ever since the advent of Herbst's argument, entropy-based methods have played an increasingly important role in the concentration literature. In fact, at the time of writing this thesis, they are probably the most active tool of research in concentration inequalities. We have already discussed the logarithmic Sobolev approach to concentration as developed by Ledoux [61] and Massart [69]. In this section, we briefly describe two other important information theoretic methods, namely, the modified log-Sobolev inequalities of Ledoux and the transportation cost inequalities of Marton.

Modified log-Sobolev inequalities. A probability measure μ on $\mathbb{R}^n$ is said to satisfy a *modified logarithmic Sobolev inequality* if there is a function $\beta(\rho) \geq 0$ on $\mathbb{R}$, such that, whenever $\|\nabla f\|_\infty \leq \rho$,

$$\text{Ent}_\mu(e^f) \leq \beta(\rho) \int \|\nabla f\|^2 e^f d\mu$$

for all locally absolutely continuous f such that $\int e^f d\mu < \infty$.

Modified log-Sobolev inequalities were introduced by Bobkov and Ledoux [16] to provide an easier alternative to Talagrand's method for proving exponential concentration. They showed that like log-Sobolev inequalities, modified log-Sobolev inequalities also tensorize in a certain sense. An important consequence of this is the following concentration result for product of measures which satisfy Poincaré inequalities (the exponential distribution being the prototype for such measures):

Theorem 2.11 *Suppose μ is a probability measure on $\mathbb{R}$ satisfying a Poincaré inequality with constant c ; that is, for any locally absolutely continuous f , $\text{Var}_\mu(f) \leq c \int |f'|^2 d\mu$. Then, any function $F : \mathbb{R}^n \rightarrow \mathbb{R}$ satisfying*

$$\sum_{i=1}^n \left| \frac{\partial F}{\partial x_i} \right|^2 \leq a^2 \quad \text{and} \quad \max_{1 \leq i \leq n} \left| \frac{\partial F}{\partial x_i} \right| \leq b$$

μ^n -almost everywhere is integrable with respect to μ^n , and for every $t \geq 0$,

$$\mu^n\{F \geq \int F d\mu^n + t\} \leq \exp\left(-\frac{1}{K} \min\left(\frac{t}{b}, \frac{t^2}{a^2}\right)\right).$$

This occurs as Corollary 5.15 in [61]. For more on modified log-Sobolev inequalities, one can look at the survey [60]. Some recent developments using the modified log-Sobolev approach (alternatively, the “entropy method”) have already been discussed in section 2.1.

Transportation cost inequalities. Transportation cost inequalities were introduced by Marton [65] as a version of measure concentration which works by investigating distances between measures.

Let’s begin with some familiar definitions. The informational divergence (or Kullback-Leibler divergence, or relative entropy) of a measure ν with respect to another measure μ on the same space is defined as:

$$D(\nu\|\mu) := \int \log\left(\frac{d\nu}{d\mu}\right) d\nu.$$

If these measures are probability measures defined on a Polish space $(\mathcal{X}, d)$, then the L^1 and L^2 Wasserstein distances between ν and μ are defined as

$$W_1(\nu, \mu) := \inf_{\pi} \mathbb{E}_{\pi} d(Y, X) \quad \text{and} \quad W_2(\nu, \mu) := \inf_{\pi} [\mathbb{E}_{\pi} d(Y, X)^2]^{1/2}$$

where Y and X are random variables distributed according to the laws ν and μ , and the infimum is taken over all distributions π on $\mathcal{X}^2$ that have ν and μ as marginals.

A probability measure μ on $\mathcal{X}$ is said to satisfy a *transportation cost inequality* with constant c , if

$$W_1(\nu, \mu) \leq \sqrt{2cD(\nu\|\mu)} \quad \text{for all probability measures } \nu \text{ on } \mathcal{X}.$$

Similarly, μ satisfies a *quadratic transportation cost inequality* (or, as Marton prefers

to call it, a distance-divergence inequality) if

$$W_2(\nu, \mu) \leq \sqrt{2cD(\nu\|\mu)} \quad \text{for all probability measures } \nu \text{ on } \mathcal{X}.$$

It is not difficult to deduce from either of the above conditions that for any two measurable sets $A, B \subseteq \mathcal{X}$,

$$d(A, B) \leq \sqrt{2c \log \frac{1}{\mu(A)}} + \sqrt{2c \log \frac{1}{\mu(B)}},$$

where $d(A, B) := \inf\{d(x, y) : x \in A, y \in B\}$. Putting $B = \{x : d(x, A) \geq t\}$ in the above, we obtain an abstract measure concentration inequality,

$$1 - \mu(A_t) \leq e^{-t^2/8c},$$

where $A_t := \{x : d(x, A) < t\}$. This interpretation of measure concentration was put forward by Milman in the late seventies. In particular, it easily implies concentration inequalities for Lipschitz functions. This will be discussed in some detail in section 2.5.

The above was the original line of argument by Marton connecting transportation cost inequalities with concentration of measure.

Quadratic transportation cost inequalities have been more successful for Euclidean spaces, mainly because they have a tensorizing property similar to log-Sobolev inequalities, which is not true for transportation cost inequalities based on the W_1 metric.

There seems to be a close connection between log-Sobolev and quadratic transportation cost inequalities. Indeed, Otto & Villani [80] proved that any measure which satisfies a logarithmic Sobolev inequality also satisfies a quadratic transportation cost inequality. The converse, however, is still an open question. For more information on mass transportation, we refer to the recent treatise [113].

One significant success of the transportation cost method has been its application to concentration for Markov chains. Originally proved by Marton [66] for contracting

Markov chains, the results were later extended to Doeblin recurrent chains and Φ -mixing processes by Samson [95] and simultaneously by Marton in a manuscript which was not sent for publication. We shall now briefly describe the result for contracting Markov chains.

Let μ be a probability measure on $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n$ induced by a Markov chain with successive transition kernels Π_i , $i = 1, \dots, n$; that is,

$$d\mu(x_1, \dots, x_n) = \Pi_1(dx_1)\Pi_2(x_1, dx_2) \cdots \Pi_n(x_{n-1}, dx_n).$$

Assume that the Markov chain is *contracting*, that is, there exists $\rho < 1$ such that for each $2 \leq i \leq n$ and $x, y \in \mathcal{X}_i$,

$$d_{TV}(\Pi_i(x, \cdot), \Pi_i(y, \cdot)) \leq \rho.$$

Then, the following analog of Theorem 2.6 holds:

Theorem 2.12 [Marton [66]] *For any measurable nonempty subset A of $\mathcal{X}$,*

$$\int e^{(1-\rho)^2 D_A(x)^2/4} d\mu(x) \leq \frac{1}{\mu(A)}.$$

As in Theorem 2.6, this implies

$$\mu\{D_A \geq t\} \leq \frac{1}{\mu(A)} e^{-(1-\rho)^2 t^2/4}.$$

Consequently, if $f : \mathcal{X} \rightarrow \mathbb{R}$ is a map with median m_f and $r(t)$ is a function such that whenever $f(x) > m_f + t$ and $f(y) \leq m_f$, there exists a vector $c(x) \in \mathbb{R}^n$ (not depending on y) with norm 1 such that $f(x) - f(y) \leq \sum_{i=1}^n c_i(x) \mathbb{I}\{x_i \neq y_i\}$, then

$$\mu\{f - m_f \geq t\} \leq 2e^{-(1-\rho)^2 r(t)^2/4}.$$

At this point, we should also mention that there is some recent work of Houdré and Tetali [54, 55] on concentration for Markov chains using a different approach, which we shall not discuss here.

Since the pioneering work of Marton, a significant number of important contributions were made by various authors. Talagrand [108] proved, among other things, that the gaussian measure on $\mathbb{R}^n$ satisfies a transportation cost inequality. Dembo [26] used the transportation cost and other information theoretic methods to rederive most of Talagrand's abstract inequalities. The papers by Bobkov & Götze [17], and Bobkov, Gentil & Ledoux [15] are also important. We refer to Ledoux [61], Chapter 6, for details.

More recently, Marton [67, 68] has worked on developing transportation cost inequalities for highly dependent systems of random variables that usually occur in statistical physics. In such models, the only tractable objects are the conditional distributions of small subcollections given the rest. The results in [67, 68] hold under Dobrushin-Shlosman type contractivity conditions. Dobrushin type conditions of weak dependence, which originated in statistical physics, will be discussed in section 4.2, where we shall also present a method based on exchangeable pairs, for directly obtaining concentration inequalities with explicit constants under similar situations.

2.5 Geometric measure concentration

This section contains a brief discussion of a closely related topic, called “concentration of measure” in geometric analytic circles. It is, in fact, a generalization of the idea of concentration inequalities, which has been a topic of interest in geometric functional analysis and convex geometry in the last three decades. The idea rests on the following basic observation: In high dimensional spaces, certain probability measures, including products of well behaved one-dimensional measures, exhibit the “concentration property”, which means that any set which has measure $\geq 1/2$, “engulfs” most of the space when slightly expanded. The idea was pioneered by Paul Lévy [63] who observed this phenomenon for the uniform measure on high dimensional spheres; but the connection of his work with concentration inequalities remained obscure for many years until the papers by Amir & Milman [3] and Gromov & Milman [47] revived mathematical interest in this very fundamental feature of high dimensional measure spaces.

Let us now formalize the notion expressed vaguely in the last paragraph. Given a Polish space $(\mathcal{X}, d)$ and a probability measure μ on $\mathcal{X}$, the “concentration function” $\alpha_{(\mathcal{X}, d, \mu)} : [0, \infty) \rightarrow [0, 1]$ is defined as:

$$\alpha_{(\mathcal{X}, d, \mu)}(t) := \sup\{1 - \mu(A_t) : A \subseteq \mathcal{X}, \mu(A) \geq 1/2\},$$

where $A_t := \{x \in \mathcal{X} : d(x, A) < t\}$, with $d(x, A) := \inf\{d(x, y) : y \in A\}$, as usual. This definition is due to Amir and Milman [3]. A measure is “concentrated” when α decreases rapidly as t grows. The connection with concentration inequalities for Lipschitz maps comes through the following easy observation: If $f : \mathcal{X} \rightarrow \mathbb{R}$ is a Lipschitz function with Lipschitz constant $\|f\|_{\text{Lip}}$, and m_f is a median of f (that is, $\mu\{f \geq m_f\} \geq 1/2$ and $\mu\{f \leq m_f\} \geq 1/2$), then

$$\mu\{|f - m_f| \geq t\} \leq 2\alpha_{(\mathcal{X}, d, \mu)}(t/\|f\|_{\text{Lip}}).$$

The advantage of using concentration functions is that they also apply to functions which are not necessarily Lipschitz, and indeed, not well-behaved in any classical sense, as is demonstrated for instance by Talagrand’s treatment of the concentration problem for the longest increasing subsequence of a random permutation ([107], section 7.1).

A major success of the geometric approach to measure concentration came with the simple proof of the famous Dvoretzky theorem by Milman [75]. To state Dvoretzky’s theorem, we first need a definition: A Banach space $(E, \|\cdot\|)$ is said to contain a subspace $(1 + \varepsilon)$ -isomorphic to $\mathbb{R}^k$ if there are vectors $v_1, \dots, v_k$ in E such that for all $t = (t_1, \dots, t_k) \in \mathbb{R}^k$,

$$(1 - \varepsilon)\|t\| \leq \left\| \sum_{i=1}^k t_i v_i \right\| \leq (1 + \varepsilon)\|t\|.$$

Dvoretzky’s theorem is the following very important result:

Theorem 2.13 [Dvoretzky’s Theorem [38]] *For each $\varepsilon > 0$ there exists $\eta(\varepsilon) > 0$ such that every Banach space E of dimension n contains a subspace $(1 + \varepsilon)$ -isomorphic to*

$\mathbb{R}^k$ where $k = \lceil \eta(\varepsilon) \log n \rceil$.

This result was at the center of the vigorous activity around the local theory of Banach spaces in the decades 1970-90. Milman's treatment and subsequent developments show that Dvoretzky's theorem can be viewed as a manifestation of the measure concentration phenomenon in a certain sense. We refer the interested reader to the lecture notes [77] for extensive details.

For further details about the functional analytic aspect (which is not relevant to this dissertation), we refer to Milman [76] and Chapters 2 and 3 of Ledoux [61].

2.6 Concentration on groups

A group G is called “topological” if it is endowed with a topology which makes the group operations continuous. It is a classical result that on any compact topological group G , there exists a unique probability measure μ which is left and right invariant; that is, if X is a G -valued random variable following the law μ , then xX and Xx have the same law as X for every $x \in G$. It also follows that X^{-1} has the same law. This measure μ is called the “Haar measure on G ” (or the “normalized Haar measure” in some texts, but Haar measures will always be normalized for us). For a self-contained proof of the existence and uniqueness of Haar measures on compact groups, we refer to Rudin [93], Theorem 5.14.

There is not much literature on the concentration of Haar measures. Before stating whatever little we could find, let us clarify that the works of Gromov & Milman [47] and Pestov [84, 85] are about a different kind of “concentration” on groups, which is not related to our investigation.

One early result is due to Maurey [70], who investigated the Haar measure on the group S_n of all permutations of n elements.

Theorem 2.14 [Maurey [70]] *Let μ denote the uniform probability measure on S_n . Let d_n denote the normalized Hamming metric on S_n : $d_n(\sigma, \tau) = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{\sigma(i) \neq \tau(i)\}$. Then, for any $A \subseteq S_n$ such that $\mu(A) \geq 1/2$, and any $t \geq 0$, we have $\mu(A_t) \geq 1 - 2e^{-nt^2/64}$, where $A_t = \{\sigma \in S_n : d_n(\sigma, A) < t\}$, as usual. Consequently,*

for any function $f : S_n \rightarrow \mathbb{R}$ which satisfies $|f(\sigma) - f(\tau)| \leq d_n(\sigma, \tau)$ for all $\sigma, \tau \in S_n$, we have

$$\mu\{|f - m_f| \geq t\} \leq 2e^{-nt^2/64} \text{ for all } t \geq 0,$$

where m_f is the median of f with respect to μ .

Maurey's result was generalized in the lecture notes of Milman and Schechtman ([77], Theorem 7.12) using the classical martingale argument to give the following theorem:

Theorem 2.15 [Milman & Schechtman [77]] *Let G be a group, compact with respect to a translation invariant metric d . Let $G = G_0 \supseteq G_1 \supseteq \dots \supseteq G_n = \{1\}$ be a decreasing sequence of closed subgroups of G . Let a_k be the diameter of G_{k-1}/G_k , $k = 1, \dots, n$. Let μ be the Haar measure and let $f : G \rightarrow \mathbb{R}$ be a function satisfying $|f(x) - f(y)| \leq d(x, y)$ for all $x, y \in G$. Then for all $t \geq 0$,*

$$\mu\{|f - \int f d\mu| \geq t\} \leq 2 \exp\left(-\frac{t^2}{4 \sum_{k=1}^n a_k^2}\right).$$

Moreover, if $A \subseteq G$ is such that $\mu(A) \geq 1/2$, then for all $t > 0$,

$$\mu(A_t) \geq 1 - 2 \exp\left(-\frac{t^2}{16 \sum_{k=1}^n a_k^2}\right),$$

where $A_t := \{x \in G : d(x, A) < t\}$.

Maurey's theorem may be easily recovered by considering the tower $S_n \supseteq S_{n-1} \supseteq \dots \supseteq S_0 = \{1\}$. Although the theorem looks pretty general, it is not very clear what sorts of examples it might cover. In fact, we could not find in the literature any interesting application of this result other than the original application to rederive Maurey's theorem. Still, we decided to include it in the review because it has the looks of a powerful general result which has not yet been fully exploited.

Next, let us discuss a result of Talagrand about the concentration of the Haar measure on S_n from his seminal work [107]. For each $A \subseteq S_n$ and $\sigma \in S_n$, define

$$U_A(\sigma) := \{s \in \{0, 1\}^n : \text{for some } \tau \in A, \mathbb{I}\{\tau(i) \neq \sigma(i)\} \leq s_i \text{ for each } i\}.$$

Let $V_A(\sigma)$ be the convex hull of $U_A(\sigma)$ in $[0, 1]^n$, and let

$$f(A, \sigma) := \inf\{\|s\|^2 : s \in V_A(\sigma)\}.$$

Then, Talagrand ([107], Theorem 5.1) has the following result:

Theorem 2.16 [Talagrand [107]] *For every $A \subseteq S_n$, we have*

$$\int_{S_n} e^{f(A, \sigma)/16} d\mu(\sigma) \leq \frac{1}{\mu(A)},$$

where μ is the uniform probability on S_n .

As the reader might have observed, it is not clear what is going on. Talagrand does not provide any example for the above theorem in his paper, although applications should be quite similar to those for product measures, because the setup is more or less the same. A workable corollary of this theorem was devised by McDiarmid [72].

Finally, let us state a result from Gromov & Milman [47] about the concentration of the Haar measure on SO_n — the group of $n \times n$ orthogonal matrices with determinant 1.

Theorem 2.17 [Gromov & Milman [47]] *Consider the group SO_n of $n \times n$ orthogonal matrices with determinant 1. Let d be the Hilbert-Schmidt metric on SO_n , defined as $d(A, B) = [\sum_{i,j} (a_{ij} - b_{ij})^2]^{1/2}$, where $A = (a_{ij})$ and $B = (b_{ij})$ are any two elements of SO_n . Let μ be the Haar measure on SO_n . Then, for any set $A \subseteq SO_n$ such that $\mu(A) \geq 1/2$, and for any $t \geq 0$, we have*

$$\mu(A_t) \geq 1 - \sqrt{\frac{\pi}{8}} e^{-(n-1)t^2/8}.$$

In particular, if $f : SO_n \rightarrow \mathbb{R}$ is a function such that $|f(A) - f(B)| \leq d(A, B)$ for all A, B , then $\mu\{|f - m_f| \geq t\} \leq \sqrt{\frac{\pi}{2}} e^{-(n-1)t^2/8}$ for each $t \geq 0$, where m_f is the median of f with respect to μ .

This result was used by Voiculescu [114] in his seminal work connecting free probability theory to random matrices. However, it is not of much use for a quantitative

analysis of the eigenvalue spectrum. In section 4.6, we shall develop a new method, using the *rank metric*, defined as $d(M, N) = \text{rank}(M - N)$, for analyzing the concentration of unitary matrices which will be suitable for studying spectral distributions of related random matrices which arise in free probability theory.

2.7 Stein's method of exchangeable pairs

Stein's method was introduced by Charles Stein [101] in the context of normal approximation for sums of dependent random variables. Stein's version of his method, best described as the “method of exchangeable pairs”, attained maturity in his later work [102]. A reasonably large literature has developed around the subject, but it has almost exclusively developed as a method of proving distributional convergence with error bounds. Stein's attempts at getting large deviations in [102] did not, unfortunately, prove fruitful. The main purpose of this dissertation is to outline a simple way of deriving concentration inequalities using the method of exchangeable pairs, and applying it to problems involving dependent variables.

We shall now briefly describe Stein's method for distributional approximation. Suppose we want to show that a random variable X taking values in some space $\mathcal{X}$ has approximately the same distribution as some other random variable Z . The procedure involves four steps:

1. Identify a “characterizing operator” T for Z , which has the defining property that for any function g belonging to a fixed large class of functions, $\mathbb{E}Tg(Z) = 0$. For instance, if $\mathcal{X} = \mathbb{R}$ and Z is a standard gaussian random variable, then $Tg(x) := g'(x) - xg(x)$ is a characterizing operator, acting on all locally absolutely continuous g with moderate growth at infinity.
2. Construct a random variable X' such that (X, X') is an exchangeable pair.
3. Find an operator α such that for any suitable $h : \mathcal{X} \rightarrow \mathbb{R}$, αh is an antisymmetric function (that is, $\alpha h(x, y) \equiv -\alpha h(y, x)$) and

$$|\mathbb{E}(\alpha h(X, X') | X = x) - Th(x)| \leq \varepsilon_h,$$

where ε_h is a small error depending only on h .

4. Take a function g and find h such that $Th(x) = g(x) - \mathbb{E}g(Z)$. By antisymmetry of αh , it follows that $\mathbb{E}(\alpha h(X, X')) = 0$. Combining with the previous step, we get $|\mathbb{E}g(X) - \mathbb{E}g(Z)| \leq \varepsilon_h$.

Note that the operator S defined as

$$Sh(x) := \mathbb{E}(\alpha h(X, X') | X = x)$$

is a characterizing operator for the distribution of X . Thus, the basic principle of Stein's method is to prove the closeness of the distributions of X and Z by showing that the characterizing operators are close.

Stated differently, this can be viewed as the study of stationary distributions of reversible Markov chains using their generators, but differences exist. The essential difference is that Stein's method involves only the pair (X, X') instead of the whole chain. The restriction of attention makes a lot of conceptual and practical difference.

There are other variants of Stein's method, most notably the dependency graph approach popularised by Arratia, Goldstein and Gordon [6], and the size-biased and zero-biased couplings of Barbour, Holst & Janson [10], but we shall not discuss those.

Stein's method has been successfully used to prove convergence to gaussian and Poisson distributions in various situations involving dependent random variables. (Poisson approximation by Stein's method was introduced by Chen [25] and became popular after the publication of [5, 6].) It is not our purpose here to go deeply into the regular versions of Stein's method. For further references and exposition, we refer to the recent monograph [30]. For applications of the method of exchangeable pairs and other versions of Stein's method to Poisson approximation, one can look at the survey paper by Chatterjee, Diaconis & Meckes [24].

Chapter 3

Theory and examples: Part I

In this chapter, we shall concentrate on the “abstract” part of our theory. The more directly applicable part will be presented in the next chapter.

Each theorem will be followed by a demonstrative example, which will always be the easiest application, involving sums of independent random variables. Let us now briefly mention the examples that we shall work out in this chapter.

In section 3.3, we shall obtain a simple and explicit concentration bound for $m - \tanh(\beta m)$, where m is the magnetization in the Curie-Weiss model of ferromagnetic interaction (which will be defined in that section for the reader who is not familiar with ferromagnetic models). By a generalization of this example, we shall derive in section 3.4 a broad condition under which a mathematically dubious but often useful technique from physics, known as *the naive mean field equations*, is valid. In section 3.5, we shall work out an example about estimation in models of dependent data (like Markov Random Field models) from mathematical statistics. In section 3.7, we shall apply our techniques to obtain tail bounds of the correct order for (a) the number of fixed points in a random permutation, and (b) the Spearman’s footrule distance between a random permutation and the identity. Finally, in section 3.10, we shall obtain a temperature-free concentration result about the Sherrington-Kirkpatrick model of spin glasses.

Throughout the remainder of this thesis, unless otherwise specified, the following notation will remain fixed and understood without mention:

- $\mathcal{X}$ is a Polish space and X is a random variable taking values in $\mathcal{X}$.
- $f : \mathcal{X} \rightarrow \mathbb{R}$ is a measurable map. The object of interest is $f(X)$, and we assume, without loss of generality, that $\mathbb{E}f(X) = 0$. (We shall abandon this last assumption on certain occasions.)
- X' is another random variable defined on the same probability space as X , such that (X, X') is an exchangeable pair.
- $F : \mathcal{X}^2 \rightarrow \mathbb{R}$ is a measurable map such that $F(X, X') = -F(X', X)$ almost surely (i.e. F is *antisymmetric*) and $\mathbb{E}(F(X, X')|X) = f(X)$. For now, we shall assume that both F and f are known. In Chapter 4, we shall investigate a method for obtaining F from a given f using couplings.
- We associate a function v with f , which will serve as a seemingly crude but useful “stochastic bound on the size of $f(X)^2$ ”:

$$v(x) := \frac{1}{2} \mathbb{E}(|(f(X) - f(X'))F(X, X')| | X = x). \quad (3.1)$$

The general principle of Stein’s method of exchangeable pairs is to study the behavior of $f(X)$ using $F(X, X')$. Exchangeable pairs often have natural constructions, and the theory usually gives us ways to infer about $f(X)$ if we know things about $F(X, X')$; however, getting information about F is usually the difficult part in practice. As mentioned before, we shall deal with that in Chapter 4.

In the next two sections we are going to present two basic theoretical results. We shall work out our first serious example in section 3.3.

3.1 A variance formula

In this section, we establish a formula for the variance of $f(X)$. The importance of this simple formula lies in the fact that in a sense, it contains the essence of our theory; further embellishments are technical.

We begin with the following fundamental lemma, which is the foundation of all further investigation:

Lemma 3.1 *For any measurable $h : \mathcal{X} \rightarrow \mathbb{R}$ such that $\mathbb{E}|h(X)F(X, X')| < \infty$, we have*

$$\mathbb{E}(h(X)f(X)) = \frac{1}{2}\mathbb{E}((h(X) - h(X'))F(X, X')).$$

Proof. Note that $\mathbb{E}(h(X)f(X)) = \mathbb{E}(h(X)F(X, X'))$. Using the exchangeability of X and X' , and the antisymmetric nature of F , we have

$$\mathbb{E}(h(X)F(X, X')) = \mathbb{E}(h(X')F(X', X)) = -\mathbb{E}(h(X')F(X, X')).$$

Thus,

$$\mathbb{E}(h(X)F(X, X')) = \frac{1}{2}\mathbb{E}((h(X) - h(X'))F(X, X')).$$

This completes the proof. $\square$

An immediate consequence of the above lemma is the following identity, which may be viewed as a weak law of large numbers for exchangeable pairs:

Theorem 3.2 *With the same notation as above, we have*

$$\text{Var}(f(X)) = \frac{1}{2}\mathbb{E}((f(X) - f(X'))F(X, X'))$$

whenever $\mathbb{E}(f(X)^2) < \infty$.

Proof. Recall that $\mathbb{E}(f(X)) = 0$ by assumption. Thus, we can directly use Lemma 3.1 with $h = f$. $\square$

Remark. At this point, we should mention that Reinert [89] has a technique for proving weak law type results for process valued functions using Stein's method, which she used to prove weak convergence of certain types of empirical processes. However, Reinert takes the usual Stein's method approach of treating the concentration problem as a problem of distributional approximation, where the target distribution is a point mass. Proceeding along this route cannot give exponential tail bounds.

As a second remark, note that from the above formula, we easily get $\mathbb{E}(f(X)^2) \leq \mathbb{E}(v(X))$, where v is as defined in (3.1).

Example. To quickly see how this works, let $X = \sum_{i=1}^n Y_i$, where Y_i 's are independent square integrable random variables. Let $\mu_i = \mathbb{E}(Y_i)$ and $\sigma_i^2 = \text{Var}(Y_i)$. An exchangeable pair is created by choosing a coordinate I uniformly at random from $\{1, \dots, n\}$, and defining

$$X' = \sum_{j \neq I} Y_j + Y'_I,$$

where $Y'_1, \dots, Y'_n$ are independent copies of $Y_1, \dots, Y_n$. Let $F(x, y) = n(x - y)$. Then

$$\mathbb{E}(F(X, X') | Y_1, \dots, Y_n) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(n(Y_i - Y'_i) | Y_1, \dots, Y_n) = \sum_{i=1}^n (Y_i - \mu_i).$$

Thus, we have $f(x) = x - \sum \mu_i$. The theorem now gives

$$\text{Var}(X) = \frac{n}{2} \mathbb{E}((X - X')^2) = \frac{1}{2} \sum_{i=1}^n \mathbb{E}(Y_i - Y'_i)^2 = \sum_{i=1}^n \sigma_i^2.$$

Note that this example was meant to be just a basic illustration. It requires independence of the Y_i 's, while for the variance formula to hold, we just need them to be uncorrelated. Substantial examples will be provided in sections 3.3 and 3.4.

3.2 An exponential inequality

Before working on further examples, let us describe our first concentration result, which is an exchangeable pairs version of the classical Hoeffding inequality (Theorem 2.1). Also, from now on we shall assume that

$$\mathbb{E}(e^{\theta f(X)} | F(X, X')) < \infty \quad \text{for each } \theta \in \mathbb{R} \quad (3.2)$$

in all our theorems, with the exception of Theorem 3.14, which is designed for situations where the above condition does not hold. The validity of this assumption will be trivial to check in most of our examples.

Theorem 3.3 *If C is a constant such that $|v(X)| \leq C$ almost surely (where $v(X)$ is defined in (3.1)), then we have $\mathbb{E}(e^{\theta f(X)}) \leq e^{C\theta^2/2}$ for all θ , and consequently, $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-t^2/2C}$ for each $t \geq 0$.*

Remark. A version of this result about reversible Markov kernels was observed by Schmuckenschläger [98]. (Note that an exchangeable pair is formally equivalent to a reversible kernel.) However, Schmuckenschläger restricts himself to a very special class of kernels (those with “positive curvature” in the sense of Bakry and Émery [9]) and the resulting technique becomes almost unusable in most practical problems. Indeed, the paper [98], though imaginative in many ways, has few concrete applications.

Example. To see that Theorem 3.3 is a generalization of Hoeffding’s inequality, consider $X = \sum_{i=1}^n Y_i$, where Y_i ’s are now independent random variables with $\mathbb{E}(Y_i) = \mu_i$ and $|Y_i - \mu_i| \leq c_i$ almost surely, where μ_i ’s and c_i ’s are finite constants. Creating the exchangeable pair as in the example following Theorem 3.2, and putting $f(X) = X - \mathbb{E}(X)$, we get

$$\begin{aligned} v(X) &= \frac{1}{2n} \sum_{i=1}^n \mathbb{E}(n(Y'_i - Y_i)^2 | X) \\ &= \frac{1}{2} \sum_{i=1}^n (\mathbb{E}(Y_i - \mu_i)^2 + \mathbb{E}((Y_i - \mu_i)^2 | X)) \leq \sum_{i=1}^n c_i^2, \end{aligned}$$

and so our theorem gives the usual Hoeffding bound (Theorem 2.1 in Chapter 2) in this case.

Proof of Theorem 3.3. Let $m(\theta) := \mathbb{E}(e^{\theta f(X)})$ be the moment generating function of $f(X)$. We can differentiate $m(\theta)$ and move the derivative inside the expectation because of the integrability assumption (3.2). Thus, by Lemma 3.1, we have

$$\begin{aligned} m'(\theta) &= \mathbb{E}(e^{\theta f(X)} f(X)) \\ &= \frac{1}{2} \mathbb{E}((e^{\theta f(X)} - e^{\theta f(X')}) F(X, X')). \end{aligned}$$

Now note that for any $x, y \in \mathbb{R}$,

$$\begin{aligned} \left| \frac{e^x - e^y}{x - y} \right| &= \int_0^1 e^{tx + (1-t)y} dt \\ &\leq \int_0^1 (te^x + (1-t)e^y) dt \quad (\text{by the convexity of } u \mapsto e^u) \\ &= \frac{1}{2}(e^x + e^y). \end{aligned} \tag{3.3}$$

Using this inequality, and the exchangeability of X and X' , we get

$$\begin{aligned} |m'(\theta)| &\leq \frac{|\theta|}{4} \mathbb{E}((e^{\theta f(X)} + e^{\theta f(X')})|(f(X) - f(X'))F(X, X')|) \\ &\leq \frac{|\theta|}{2} \mathbb{E}(e^{\theta f(X)}v(X) + e^{\theta f(X')}v(X')) \\ &= |\theta| \mathbb{E}(e^{\theta f(X)}v(X)) \leq C|\theta|m(\theta). \end{aligned}$$

For $\theta \geq 0$, this means $\frac{d}{d\theta} \log m(\theta) \leq C\theta$. Since $m(0) = 1$, we have $\log m(\theta) \leq C\theta^2/2$. Given $t \geq 0$, we can choose $\theta = t/C$ and get

$$\mathbb{P}\{f(X) \geq t\} \leq e^{-\theta t + \log m(\theta)} \leq e^{-t^2/2C}.$$

Similarly, $\mathbb{P}\{f(X) \leq -t\} \leq e^{-t^2/2C}$. □

3.3 Example: Curie-Weiss model

In this section, we shall work out concentration bounds for the magnetization in the Curie-Weiss model of ferromagnetic interaction. This is arguably the simplest statistical mechanical model of spin systems. For a detailed mathematical treatment of this model, we refer to the book [41] by Richard Ellis. In the next section, we shall extend our technique for the Curie-Weiss model to a more general class of models with quadratic interaction.

The state space is $\{-1, 1\}^n$, the space of all possible spin configurations of n

particles. A typical configuration will be denoted by $\sigma = (\sigma_1, \dots, \sigma_n)$. The Curie-Weiss model at inverse temperature β and a fixed external magnetic field h prescribes a joint probability density (the Gibbs measure) for these spins by the formula $p_{\beta,h}(\sigma) = Z_{\beta,h}^{-1} e^{-\beta H_h(\sigma)}$ where $Z_{\beta,h}$ is the normalizing constant, and

$$H_h(\sigma) := -\frac{1}{n} \sum_{i < j} \sigma_i \sigma_j - h \sum_{i=1}^n \sigma_i$$

is the Hamiltonian for the system. We shall henceforth assume that β and h are fixed and omit the subscripts. One quantity of interest is the magnetization, defined as

$$m = m(\sigma) := \frac{1}{n} \sum_{i=1}^n \sigma_i.$$

It is common knowledge in the physics circles (and proved rigorously in [41]) that if n is large and σ is drawn from the Gibbs measure, the random variable $m(\sigma)$ is concentrated in a neighborhood of the set of solutions of the equation

$$x = \tanh(\beta x + \beta h).$$

The equation has a unique root for small values of β (the “high temperature phase”) and multiple solutions for β above a critical range (the “low temperature phase”). For example, when $h = 0$, $\beta_c = 1$ is the critical value.

In this context, we can use Theorems 3.2 and 3.3 quite easily to prove the following result:

Proposition 3.4 *The magnetization m in the Curie-Weiss model satisfies*

$$\mathbb{E}(m - \tanh(\beta m + \beta h))^2 \leq \frac{2 + 2\beta}{n} + \frac{\beta^2}{n^2}.$$

Moreover, we also have the concentration bound

$$\mathbb{P}\left\{|m - \tanh(\beta m + \beta h)| \geq \frac{\beta}{n} + t\right\} \leq 2e^{-nt^2/(4+4\beta)}.$$

Here $\mathbb{E}$ and $\mathbb{P}$ denote expectation and probability under the Gibbs measure at inverse temperature β and external field h .

Remark. Although the Curie-Weiss model is considered to be the simplest model of ferromagnetic interaction, we haven't encountered any result in the literature which gives bounds like the above. In fact, it is unlikely that there exists a short and simple bare-hands argument which gives such bounds.

The classical way to solve the Curie-Weiss model uses ideas from large deviations theory, and involves a variational problem in the limit (see Ellis [41], Chapter IV). The magnetization is obtained by the usual technique of differentiating the free energy function. Although the limiting result is straightforward, getting finite sample tail bounds will not be easy along this route.

Proof. Suppose σ is drawn from the Gibbs distribution. We construct σ' by taking a step in the Gibbs sampler as follows: Choose a coordinate I uniformly at random, and replace the I^{th} coordinate of σ by an element drawn from the conditional distribution of the I^{th} coordinate given the rest. It is well-known and easy to prove that (σ, σ') is an exchangeable pair. Let

$$F(\sigma, \sigma') := \sum_{i=1}^n (\sigma_i - \sigma'_i).$$

Now define

$$m_i(\sigma) := \frac{1}{n} \sum_{j \leq n, j \neq i} \sigma_j, \quad i = 1, \dots, n.$$

Since the Hamiltonian is a simple explicit function, the conditional distribution of the i^{th} coordinate given the rest is easy to obtain. An easy computation gives $\mathbb{E}(\sigma_i | \{\sigma_j, j \neq i\}) = \tanh(\beta m_i + \beta h)$. Thus, we have

$$\begin{aligned} f(\sigma) &= \mathbb{E}(F(\sigma, \sigma') | \sigma) = \frac{1}{n} \sum_{i=1}^n (\sigma_i - \mathbb{E}(\sigma_i | \{\sigma_j, j \neq i\})) \\ &= m - \frac{1}{n} \sum_{i=1}^n \tanh(\beta m_i + \beta h). \end{aligned}$$

Now note that $|F(\sigma, \sigma')| \leq 2$, because σ and σ' differ at only one coordinate. Also, since the map $x \mapsto \tanh x$ is 1-Lipschitz, we have

$$|f(\sigma) - f(\sigma')| \leq |m(\sigma) - m(\sigma')| + \frac{\beta}{n} \sum_{i=1}^n |m_i(\sigma) - m_i(\sigma')| \leq \frac{2(1+\beta)}{n}.$$

Thus, by Theorem 3.2 we have

$$\text{Var}\left(m - \frac{1}{n} \sum_{i=1}^n \tanh(\beta m_i + \beta h)\right) \leq \frac{2(1+\beta)}{n}$$

and from Theorem 3.3,

$$\mathbb{P}\left\{\left|m - \frac{1}{n} \sum_{i=1}^n \tanh(\beta m_i + \beta h)\right| \geq t\right\} \leq 2e^{-nt^2/(4+4\beta)}.$$

Finally note that for each i , by the Lipschitz nature of the $\tanh$ function, we get

$$|\tanh(\beta m_i + \beta h) - \tanh(\beta m + \beta h)| \leq \beta |m_i - m| \leq \frac{\beta}{n}.$$

This completes the proof of the theorem. $\square$

3.4 Validity of naive mean field equations

The proof of Proposition 3.4 shows that there is no reason why the argument will not generalize to the Hamiltonian

$$H(\sigma) := - \sum_{i < j} J_{ij} \sigma_i \sigma_j - \sum_{i=1}^n h_i \sigma_i \tag{3.4}$$

where $J = (J_{ij})_{1 \leq i, j \leq n}$ is a symmetric interaction matrix with zeros on the diagonal, and $h_1, \dots, h_n$ are fixed real numbers. In this section, we are going to carry through an extension of our treatment of the Curie-Weiss model to find conditions under which a certain widely used (but mathematically dubitable) technique from physics — the

naive mean field equations — is valid.

Throughout this section, we shall use the standard physical notation $\langle \cdot \rangle$ to denote the expectation with respect to the Gibbs measure.

Now, it is an easy fact that $\mathbb{E}_\beta(\sigma_i | \{\sigma_j, j \neq i\}) = \tanh(\beta \sum_{j=1}^n J_{ij} \sigma_j + h_i)$. This gives what are called the *Callen equations* (see e.g. Chapter 3 of [82]):

$$\langle \sigma_i \rangle = \left\langle \tanh \left(\beta \sum_{j=1}^n J_{ij} \sigma_j + \beta h_i \right) \right\rangle, \quad i = 1, \dots, n. \quad (3.5)$$

These equations are somehow just a restatement of the model, and not useful for anything deeper. The *naive mean field equations* are a modification of the Callen equations. The general physical intuition is that, when the spins are sufficiently “uncorrelated”, the expectation on the right hand side of the Callen equations can be taken inside the tanh. This gives

$$“\langle \sigma_i \rangle = \tanh \left(\beta \sum_{j=1}^n J_{ij} \langle \sigma_j \rangle + \beta h_i \right), \quad i = 1, \dots, n”. \quad (3.6)$$

The reason why this is written within quotes is that no reasonable general condition is known for the validity of these equations. Physicists know that these are *not* valid in models with short range interactions like the Ising model, but generally work when the interactions are of very long range (like the Curie-Weiss, for instance). Rigorous texts like [41] prefer to avoid talking about mean field equations. Still, this technique is used with reasonable measures of success in a variety of fields, including computer science, image processing, neural networks and other computational sciences. A system of n equations may seem strange at first sight, but the fact is that they often reduce to a much smaller system in practice (for instance, in the Curie-Weiss model, they reduce to just one equation). For a recent survey of applications in areas outside of physics, one can look at [81]. For the physical perspective, a classical text is Parisi’s book [82].

We shall now show that, although it is difficult to obtain general conditions under which the average spins $\{\langle \sigma_i \rangle, i = 1, \dots, n\}$ satisfy the mean field equations, the

conditional averages of the spins, defined as

$$\langle \sigma_i \rangle^- := \mathbb{E}(\sigma_i | \{\sigma_j, j \neq i\}) = \tanh(\beta \sum_j J_{ij} \sigma_j + \beta h_i), \quad i = 1, \dots, n \quad (3.7)$$

satisfy (3.6) with high probability whenever the Hilbert-Schmidt norm of J (denoted by $\|J\|_{HS}$) is not too large. The precise condition is

$$\|J\|_{HS} := [\sum_{i,j} J_{ij}^2]^{1/2} \ll n^{1/3}. \quad (3.8)$$

We shall also show that the mean field equations (3.6) themselves are approximately valid at sufficiently high temperature in the class of models satisfying (3.8). The results are summarized in the following theorem:

Theorem 3.5 *Fix $\beta \geq 0$. Let $\rho = \|J\|_{HS} = [\sum_{i,j} J_{ij}^2]^{1/2}$. Let $\langle \sigma_i \rangle^-$, $i = 1, \dots, n$ be defined as in (3.7). For $1 \leq i \leq n$, define*

$$\begin{aligned} \tilde{\varepsilon}_i &:= \langle \sigma_i \rangle^- - \tanh\left(\beta \sum_{j=1}^n J_{ij} \langle \sigma_j \rangle^- + \beta h_i\right), \quad \text{and} \\ \varepsilon_i &:= \langle \sigma_i \rangle - \tanh\left(\beta \sum_{j=1}^n J_{ij} \langle \sigma_j \rangle + \beta h_i\right). \end{aligned}$$

Then we have the bound $\langle \tilde{\varepsilon}_i^2 \rangle \leq 2\beta^2(1 + \beta\rho) \sum_{j=1}^n J_{ij}^2$ for each i , and hence

$$\frac{1}{n} \sum_{i=1}^n \langle \tilde{\varepsilon}_i^2 \rangle \leq \frac{1}{n} 2(1 + \beta\rho) \beta^2 \rho^2. \quad (3.9)$$

We also have the tail bounds

$$\langle |\tilde{\varepsilon}_i| \geq t \rangle \leq 2 \exp\left(-\frac{t^2}{4\beta^2(1 + \beta\rho) \sum_{j=1}^n J_{ij}^2}\right) \quad (3.10)$$

for $1 \leq i \leq n$. Moreover, if $\beta < 1/\rho$, we have

$$\frac{1}{n} \sum_{i=1}^n \varepsilon_i^2 \leq \frac{2(1 + \beta\rho)\beta^2\rho^2}{n(1 - \beta\rho)^2}. \quad (3.11)$$

Discussion and examples. Before we come to a critical discussion, let us, for a moment, gloat over the fact that it is difficult to imagine that such bounds (even the second moment bounds) can be easily obtained by bare-hands arguments, or by any available technique, for that matter!

The first part of the theorem says that under (3.8), the conditional averages $\{\langle \sigma_i \rangle^-, i = 1, \dots, n\}$, instead of the unconditional ones, satisfy the mean field equations (3.6). In other words, we have a set of modified mean field equations

$$\langle \sigma_i \rangle^- = \tanh \left(\beta \sum_{j=1}^n J_{ij} \langle \sigma_j \rangle^- + \beta h_i \right), \quad i = 1, \dots, n, \quad (3.12)$$

which are approximately valid with high probability. In particular, if each $\langle \sigma_j \rangle^-$ is concentrated around its mean, then we can recover the mean field equations from the information in (3.9). One condition under which this happens is the “high temperature condition” $\beta < 1/\rho$, which gives the second part of the theorem. But presumably the concentration of $\langle \sigma_j \rangle^-$ can happen under other conditions also.

The condition (3.8) can be interpreted as some sort of a condition of long range interaction, as will be made clear by the following example. Consider a graph $G = (V, E)$ on $V = \{1, \dots, n\}$ with constant degree r . The (normalized) Ising Hamiltonian on $\{-1, 1\}^V$ is defined as

$$H(\sigma) = -\frac{1}{r} \sum_{(i,j) \in E} \sigma_i \sigma_j - h \sum_{i=1}^n \sigma_i,$$

where h is the external field. Thus, $J_{ij} = 1/r$ if $(i, j) \in E$, and $= 0$ otherwise. Since the graph has constant degree r , a simple computation gives $\|J\|_{HS} = \sqrt{n/r}$. Thus, the modified mean field equations (3.12) hold whenever $r \gg n^{1/3}$, which is a long range condition.

For another example, consider the following generalization of the Curie-Weiss model: Suppose each J_{ij} is bounded in magnitude by $1/n$. Then $\|J\|_{HS} \leq 1$, and therefore the modified mean field equations hold. Moreover, the second part of the theorem shows that in this class of models, if $\beta < 1$, then the mean field equations (3.6) are valid.

Proof of Theorem 3.5. Throughout this proof, we shall use the fact that $\tanh$ is a Lipschitz function without explicit mention. Also, as a natural extension of the notation in the previous section, we shall write

$$m_i = m_i(\sigma) := \sum_{j=1}^n J_{ij} \sigma_j + h_i.$$

Note that $\langle \sigma_i \rangle^- = \tanh(\beta m_i)$.

Now fix i , $1 \leq i \leq n$. Construct (σ, σ') as in the proof of Proposition 3.4, by choosing σ from the Gibbs measure and taking a step in the Gibbs sampler to get σ' as follows: Choose a coordinate I uniformly at random, and replace the I^{th} coordinate of σ by a random sample from the conditional distribution of σ_I given $\{\sigma_j, j \neq I\}$. Define $F(\sigma, \sigma') = n \sum_{j=1}^n J_{ij}(\sigma_j - \sigma'_j) = n J_{iI}(\sigma_I - \sigma'_I)$. Then, as before,

$$f(\sigma) = \sum_{j=1}^n J_{ij}(\sigma_j - \tanh(\beta m_j)) = m_i - \sum_{j=1}^n J_{ij} \tanh(\beta m_j) - h_i. \quad (3.13)$$

Now, if σ and σ' differ at site I , then

$$\begin{aligned} |f(\sigma) - f(\sigma')| &\leq 2|J_{iI}| + \sum_{j=1}^n |J_{ij}(\tanh(\beta m_j(\sigma)) - \tanh(\beta m_j(\sigma')))| \\ &\leq 2|J_{iI}| + \sum_{j=1}^n |J_{ij}| |\beta(m_j(\sigma) - m_j(\sigma'))| \\ &= 2|J_{iI}| + 2\beta \sum_{j=1}^n |J_{ij} J_{jI}|. \end{aligned}$$

Also, $|F(\sigma, \sigma^I)| \leq 2n|J_{iI}|$. Thus, we have

$$\begin{aligned}
v(\sigma) &= \frac{1}{2} \mathbb{E}(|(f(\sigma) - f(\sigma'))F(\sigma, \sigma')||\sigma) \\
&\leq \frac{1}{2} \sum_{k=1}^n (2|J_{ik}| + 2\beta \sum_{j=1}^n |J_{ij}J_{jk}|) 2|J_{ik}| \\
&\leq 2 \sum_{k=1}^n J_{ik}^2 + 2\beta \left[\sum_{j,k} J_{ij}^2 J_{ik}^2 \right]^{1/2} \left[\sum_{j,k} J_{jk}^2 \right]^{1/2} \\
&= 2(1 + \beta\rho) \sum_{j=1}^n J_{ij}^2.
\end{aligned} \tag{3.14}$$

Thus, by Theorem 3.2 and (3.13), we get

$$\left\langle \left(m_i - \sum_{j=1}^n J_{ij} \tanh(\beta m_j) - h_i \right)^2 \right\rangle = \langle f(\sigma)^2 \rangle \leq 2(1 + \beta\rho) \sum_{j=1}^n J_{ij}^2. \tag{3.15}$$

To complete the proof of the first set of inequalities in Theorem 3.5, observe that

$$\begin{aligned}
\langle \tilde{\varepsilon}_i^2 \rangle &= \left\langle \left(\tanh(\beta m_i) - \tanh \left(\beta \sum_{j=1}^n J_{ij} \tanh(\beta m_j) + \beta h_i \right) \right)^2 \right\rangle \\
&\leq \beta^2 \left\langle \left(m_i - \sum_{j=1}^n J_{ij} \tanh(\beta m_j) - h_i \right)^2 \right\rangle,
\end{aligned}$$

and combine with (3.15). The tail bound on $\tilde{\varepsilon}_i$ follows from the same analysis, using Theorem 3.3.

Next, for each i define the function $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ as $g_i(x) = \sum_{j=1}^n J_{ij} \tanh(\beta x_j) + h_i$. Define $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ as $g(x) := (g_1(x), \dots, g_n(x))$. Then note that for any $x, y \in \mathbb{R}^n$,

$$\begin{aligned}
\|g(x) - g(y)\|^2 &= \sum_{i=1}^n \left(\sum_{j=1}^n J_{ij} (\tanh(\beta x_j) - \tanh(\beta y_j)) \right)^2 \\
&\leq \beta^2 \sum_{i=1}^n \left(\sum_{j=1}^n |J_{ij}(x_j - y_j)| \right)^2 \\
&\leq \beta^2 \rho^2 \|x - y\|^2.
\end{aligned}$$

Thus, if $\beta < 1/\rho$, then g is a contraction map with respect to the Euclidean metric. Therefore by the well-known theorem about contraction maps (see, e.g. Theorem 9.23 in Rudin [94]), g has a unique fixed point, which we shall call x^* . Now note that for any $x \in \mathbb{R}^n$,

$$\begin{aligned} \|x - x^*\| &\leq \|(x - g(x)) - (x^* - g(x^*))\| + \|g(x) - g(x^*)\| \\ &\leq \|x - g(x)\| + \beta\rho\|x - x^*\|. \end{aligned}$$

Thus for any x , we have $\|x - x^*\| \leq (1 - \beta\rho)^{-1}\|x - g(x)\|$. Applying to $x = (m_1, \dots, m_n)$, and remembering that $\langle (m_i - a)^2 \rangle$ is minimized at $a = \langle m_i \rangle$, we get

$$\begin{aligned} \sum_{i=1}^n \langle (m_i - \langle m_i \rangle)^2 \rangle &\leq \sum_{i=1}^n \langle (m_i - x_i^*)^2 \rangle \\ &\leq \frac{1}{(1 - \beta\rho)^2} \sum_{i=1}^n \langle (m_i - \sum_{j=1}^n J_{ij} \tanh(\beta m_j) - h_i)^2 \rangle \\ &\leq \frac{2(1 + \beta\rho)\rho^2}{(1 - \beta\rho)^2} \quad \text{by inequality (3.15)}. \end{aligned}$$

Finally, observe that

$$\begin{aligned} \sum_{i=1}^n \varepsilon_i^2 &= \sum_{i=1}^n (\langle \tanh(\beta m_i) \rangle - \tanh(\beta \langle m_i \rangle))^2 \\ &\leq \beta^2 \sum_{i=1}^n \langle (m_i - \langle m_i \rangle)^2 \rangle. \end{aligned}$$

Combined with the previous step, this completes the proof. $\square$

3.5 Example: Least squares for the Ising model

Consider an undirected graph $G = (V, E)$, where $V = \{1, \dots, n\}$. Let r be the maximum degree of G . Let $X = (X_1, \dots, X_n)$ be a random element of $\{-1, 1\}^G$. The

Ising model assigns a probability distribution for X according to the formula

$$\mathbb{P}_\theta\{X = x\} = Z(\theta)^{-1} e^{-\theta \sum_{(i,j) \in E} x_i x_j} \quad (3.16)$$

where θ is an unknown parameter (the “inverse temperature”, usually denoted by β in the physics literature), and $Z(\theta)$ is the normalizing constant. The parameter space is $\Omega := [0, \infty)$. The natural statistical problem in this model is the following: How to make inference about θ from a single realization of X ? This is one of most elementary (yet analytically almost intractable) models of dependent discrete data.

The classical maximum likelihood approach for this problem has been discussed in detail by Pickard [86]. The main difficulty with directly computing the maximum likelihood estimator (MLE) is that the normalizing constant has no explicit form except in very special cases, and there is no polynomial time algorithm for exact numerical evaluation.

In a well-cited paper, Geyer & Thompson [46] devised a feasible Monte Carlo technique for computing the MLE in models like the above. One of the examples considered in that paper involves an instance of the autologistic model of Besag [12], which is a generalization of the Ising model. The autologistic model of binary data assumes that the conditional distribution of each X_i given the rest can be modeled as a logistic regression, that is,

$$\log \frac{\mathbb{P}\{X_i = 1 | (X_j, j \neq i)\}}{1 - \mathbb{P}\{X_i = 1 | (X_j, j \neq i)\}} = \alpha_i + \sum_{j \leq n, j \neq i} \beta_{ij} X_j,$$

where the α_i ’s and the β_{ij} ’s are known functions of a collection of unknown parameters. A simple verification shows that the Ising model described by (3.16) is a special case of the autologistic model.

Monte Carlo algorithms for computing the MLE in such problems are widely used nowadays, but lingering doubts remain about the rates of convergence, specially in models with high degrees of dependence like the ones described above.

A method which does not require simulations, but fell out of favor due to efficiency issues after the advent of Monte Carlo techniques, is the maximum pseudolikelihood

approach introduced by Besag [13]. Besag's pseudolikelihood function is defined as

$$f(\theta|X) := \prod_{i=1}^n f_i(\theta|X),$$

where $f_i(\theta|X)$ is the conditional density of X_i given $(X_j, j \neq i)$ under θ . Now, if X is a gaussian vector such that $\text{Var}_\theta(X_i|(X_j, j \neq i))$ is a constant independent of i and θ , the pseudolikelihood problem reduces to minimizing

$$S(\theta) := \frac{1}{n} \sum_{i=1}^n (X_i - \mathbb{E}_\theta(X_i|(X_j, j \neq i)))^2, \quad (3.17)$$

which may be called a “conditional least squares problem”. We shall now show that somewhat surprisingly, minimizing $S(\theta)$ to estimate θ may be a good idea even outside the gaussian framework. In particular, we shall show by way of example later in this section that it works in the Ising model, and the method can even be extended to construct nonasymptotic confidence intervals for θ . One of the interesting consequences is the following:

Proposition 3.6 *In the Ising model on a graph with maximum degree r as defined in (3.16), we have the bound*

$$\mathbb{E}_\theta(S(\theta) - \inf_{\psi} S(\psi)) \leq C \sqrt{\frac{r \log n}{n}} \quad \text{for every } \theta \geq 0,$$

where C is a numerical constant.

This shows that the conditional least squares inference for θ should work in this model when $r \ll n/\log n$. However, there is a caveat: Although our results will show that $S(\theta)$ is minimized near the true value of θ under pretty general conditions, they say nothing about the sharpness of the minimum of $S(\theta)$. If S is too flat, our results will be of no use.

In the remainder of this section, we shall adopt the notation $\bar{x}^i$ for the vector $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$, where x is the original vector $(x_1, \dots, x_n)$. Let $X = (X_1, \dots, X_n)$ be a random vector in $[-1, 1]^n$, whose distribution is parametrized by a

parameter θ belonging to some parameter space Ω . For $1 \leq i \leq n$, let

$$\mu_i(\theta, \bar{x}^i) := \mathbb{E}_\theta(X_i | \bar{X}^i = \bar{x}^i),$$

For each θ , let $\{a_{ij}(\theta)\}_{1 \leq i, j \leq n}$ be an array of nonnegative real numbers with zeros on the diagonal such that for any x, y, i and θ we have

$$|\mu_i(\theta, \bar{x}^i) - \mu_i(\theta, \bar{y}^i)| \leq \sum_{j=1}^n a_{ij}(\theta) \mathbb{I}\{x_j \neq y_j\},$$

and let

$$M := \frac{1}{n} \sup_{\theta \in \Omega} \sum_{i,j} a_{ij}(\theta). \quad (3.18)$$

Lemma 3.7 below will show that when $M \ll n$, which holds in many models (including Ising type models), $S(\psi) - S(\theta)$ is “approximately nonnegative with high probability” whenever θ is the true value of the parameter and ψ is any other value. This will be shown by decomposing $S(\psi) - S(\theta)$ as $A(\psi, \theta) + B(\psi, \theta)$, where $A \geq 0$ and $B \approx 0$ under $\mathbb{P}_\theta$. The explicit expressions for A and B are as follows:

$$A(\psi, \theta) = \frac{1}{n} \sum_{i=1}^n (\mu_i(\theta, \bar{X}^i) - \mu_i(\psi, \bar{X}^i))^2$$

and

$$B(\psi, \theta) = \frac{2}{n} \sum_{i=1}^n (\mu_i(\theta, \bar{X}^i) - \mu_i(\psi, \bar{X}^i))(X_i - \mu_i(\theta, \bar{X}^i)).$$

It is easy to check that indeed $S(\psi) - S(\theta) = A(\psi, \theta) + B(\psi, \theta)$. It is also apparent that $A(\psi, \theta) \geq 0$. The following lemma, the proof of which depends heavily on our techniques, shows that $B(\psi, \theta) \approx 0$ with high probability under θ whenever $M \ll n$. It is not completely honest, though, because the constants are bad.

Lemma 3.7 *Suppose X takes values in $[-1, 1]^n$ and let M , A and B be defined as above. Then for any $\psi, \theta \in \Omega$ and $t \geq 0$, we have*

$$\mathbb{P}_\theta\{|B(\psi, \theta)| \geq t\} \leq 2e^{-nt^2/(96M+32)}.$$

Proof. Fix $\psi, \theta \in \Omega$. We produce X' , as usual, by taking a step in the Gibbs sampler: A coordinate I is chosen uniformly at random, and X_I is replaced by X'_I drawn from the conditional distribution (under $\mathbb{P}_\theta$) of the I^{th} coordinate given $\bar{X}^I$. Next, for each i , let

$$g_i(\bar{x}^i) := \mu_i(\theta, \bar{x}^i) - \mu_i(\psi, \bar{x}^i),$$

and define

$$F(x, y) = \sum_{i=1}^n (g_i(\bar{x}^i) + g_i(\bar{y}^i))(x_i - y_i).$$

Clearly, F is antisymmetric. Note that

$$\begin{aligned} f(X) &:= \mathbb{E}_\theta(F(X, X')|X) = 2\mathbb{E}_\theta(g_I(\bar{X}^I)(X_I - X'_I)|X) \\ &= \frac{2}{n} \sum_{i=1}^n g_i(\bar{X}^i)(X_i - \mu_i(\theta, \bar{X}^i)). \end{aligned}$$

Thus, in our notation, $f(X) = B(\psi, \theta)$. From the given conditions it easily follows that if x and y are elements of $[-1, 1]^n$ which differ only in the i^{th} coordinate, then

$$|f(x) - f(y)| \leq \frac{4}{n} \left(\sum_{j=1}^n (a_{ji}(\psi) + 2a_{ji}(\theta)) + 1 \right).$$

Also, quite clearly, $|F(X, X')| \leq 8$. Combining, we have

$$\begin{aligned} v(X) &= \frac{1}{2} \mathbb{E}_\theta(|(f(X) - f(X'))F(X, X')||X) \\ &\leq \frac{16}{n^2} \sum_{i=1}^n \sum_{j=1}^n (a_{ji}(\psi) + 2a_{ji}(\theta)) + \frac{16}{n}. \end{aligned}$$

Invoking Theorem 3.3 completes the proof. $\square$

The above lemma is only a moral justification for using the “conditional least squares approach” for estimating parameters in models of dependent data. It does not show that the true value of θ is an approximate *global* minimizer of $S(\theta)$. For that,

and also for constructing confidence regions, we need tail bounds on $S(\theta) - \inf_{\psi} S(\psi)$. For instance, a $(1 - \alpha)$ -level *confidence region* for the true value of θ can be defined as $\{\theta : S(\theta) - \inf_{\psi \in \Omega} S(\psi) \leq t_{\alpha}\}$, where t_{α} is chosen such that for any θ ,

$$\mathbb{P}_{\theta}\{S(\theta) - \inf_{\psi} S(\psi) > t_{\alpha}\} \leq \alpha.$$

For doing all that, we need to introduce some more notation. Define a pseudometric d on Ω as follows:

$$d(\theta, \theta') := \sup_x \frac{1}{n} \sum_{i=1}^n |\mu_i(\theta, \bar{x}^i) - \mu_i(\theta', \bar{x}^i)|.$$

For every $\varepsilon > 0$, let $N_d(\varepsilon)$ denote the minimum number of closed ε -balls (w.r.t. the metric d) required to cover Ω . We have the following result:

Theorem 3.8 *Suppose X takes values in $[-1, 1]^n$. Let $S(\theta)$ and M be defined as in (3.17) and (3.18), and let N_d be the covering number defined above. Then, for any $\varepsilon > 0$ and any $\theta \in \Omega$, we have*

$$\mathbb{P}_{\theta}\{S(\theta) - \inf_{\psi \in \Omega} S(\psi) \geq 4\varepsilon + t\} \leq 2N_d(\varepsilon)e^{-nt^2/(96M+32)}$$

for all $t \geq 0$. Consequently, we also have

$$\mathbb{E}_{\theta}(S(\theta) - \inf_{\psi \in \Omega} S(\psi)) \leq \inf_{\varepsilon > 0} \left\{ 4\varepsilon + 2\sqrt{\frac{(96M+32) \log(2N_d(\varepsilon))}{n}} \right\}.$$

Application to the Ising model. For the Ising model described at the beginning of the section, it is easy to verify that $\mu_i(\theta, \bar{x}^i) = \tanh(\theta \sum_{j \in N(i)} x_j)$, where $N(i)$ is the neighborhood of i in G . Now, since $\tanh x \in [-1, 1]$ for all $x \in \mathbb{R}$, we have

$$|\mu_i(\theta, \bar{x}^i) - \mu_i(\theta, \bar{y}^i)| \leq 2 \sum_{j \in N(i)} \mathbb{I}\{x_j \neq y_j\}.$$

Thus, we can take $a_{ij}(\theta) = 2\mathbb{I}\{(i, j) \in E\}$ irrespective of θ . With this choice of a_{ij} 's, we have $M = 2r$, where recall that r is the maximum degree of G .

Now note that $\sum_{j \in N(i)} x_j \in \{0, \pm 1, \dots, \pm r\}$, and $\tanh$ is an odd function. Therefore for any θ, θ' and x ,

$$|\mu_i(\theta, \bar{x}^i) - \mu_i(\theta', \bar{x}^i)| \leq \sup_{s \in \{0, 1, \dots, r\}} |\tanh(\theta s) - \tanh(\theta' s)|.$$

Since $\tanh$ is a Lipschitz function, therefore the right hand side is bounded by $|\theta - \theta'|r$. On the other hand, since $\tanh$ is an increasing function bounded by 1, therefore the right hand side is also bounded by $1 - \tanh(\min\{\theta, \theta'\}) \leq \exp(-\min\{\theta, \theta'\})$. Combining, we have

$$d(\theta, \theta') = \sup_x \frac{1}{n} \sum_{i=1}^n |\mu_i(\theta, \bar{x}^i) - \mu_i(\theta', \bar{x}^i)| \leq \min\{|\theta - \theta'|r, e^{-\min\{\theta, \theta'\}}\}.$$

Now fix any $\varepsilon \in (0, 1)$. Let $L = -\log \varepsilon$. Equipartition the interval $[0, L]$ into $[Lr/2\varepsilon] + 1$ subintervals of length $\leq 2\varepsilon/r$ each. The above bound shows that the end points of these subintervals, together with L , form an ε -net with respect to the pseudometric d on $[0, \infty)$. Thus,

$$N_d(\varepsilon) \leq \frac{r|\log \varepsilon|}{2\varepsilon} + 2.$$

It is now easy to apply Theorem 3.8. The second bound directs us to the optimal choice of ε , which we take to be $\varepsilon = \sqrt{r/n}$. Theorem 3.8 now gives

$$\mathbb{P}_\theta\{S(\theta) - \inf_{\psi} S(\psi) \geq 4\sqrt{\frac{r}{n}} + t\} \leq 2\sqrt{rn}(\log n)e^{-nt^2/(96r+32)},$$

and

$$\mathbb{E}_\theta(S(\theta) - \inf_{\psi} S(\psi)) \leq C\sqrt{\frac{r \log n}{n}},$$

where C is a computable numerical constant. This proves Proposition 3.6.

Proof of Theorem 3.8. Fix $\varepsilon > 0$. Let $k = N_d(\varepsilon)$, and let $\theta_1, \dots, \theta_k$ be the centers of a collection of ε -balls which cover Ω . Then for every θ there exists i such

that $|S(\theta) - S(\theta_i)| \leq 4d(\theta, \theta_i) \leq 4\varepsilon$. Thus,

$$\min_{1 \leq i \leq k} S(\theta_i) - \inf_{\theta} S(\theta) \leq 4\varepsilon.$$

Therefore,

$$\begin{aligned} \mathbb{P}_{\theta}\{S(\theta) - \inf_{\psi} S(\psi) \geq 4\varepsilon + t\} \\ \leq \mathbb{P}_{\theta}\{|S(\theta) - \inf_{1 \leq i \leq k} S(\theta_i)| \geq t\} \leq \sum_{i=1}^k \mathbb{P}_{\theta}\{|S(\theta) - S(\theta_i)| \geq t\}. \end{aligned}$$

The first bound follows from this and Lemma 3.7. Now note that for any $b > 0$ and $a > \sqrt{e}$, we have

$$\begin{aligned} \int_0^{\infty} (ae^{-bt^2} \wedge 1) dt &= \sqrt{b^{-1} \log a} + \int_{\sqrt{b^{-1} \log a}}^{\infty} ae^{-bt^2} dt \\ &\leq \sqrt{b^{-1} \log a} + \int_{\sqrt{b^{-1} \log a}}^{\infty} \frac{t}{\sqrt{b^{-1} \log a}} ae^{-bt^2} dt \\ &= \sqrt{b^{-1} \log a} + \frac{1}{2b\sqrt{b^{-1} \log a}} \\ &\leq 2\sqrt{b^{-1} \log a} \quad (\text{since } a \geq \sqrt{e} \implies 2\sqrt{\log a} \geq 1/\sqrt{\log a}). \end{aligned}$$

Thus,

$$\begin{aligned} \mathbb{E}_{\theta}(S(\theta) - \inf_{\psi} S(\psi)) &\leq 4\varepsilon + \int_0^{\infty} (2N_d(\varepsilon)e^{-nt^2/(96M+32)} \wedge 1) dt \\ &\leq 4\varepsilon + 2\sqrt{\frac{(96M+32) \log(2N_d(\varepsilon))}{n}}. \end{aligned}$$

This completes the proof of Theorem 3.8. □

3.6 An inequality for self-bounded functions

Theorem 3.3 can be extended in several ways. The following extension is analogous to existing results for the so called “self-bounded” functions. The terminology was

introduced by Boucheron, Lugosi & Massart in [18]. Recall the notation of these authors that was described in section 2.1 of Chapter 2. They call a function “self-bounded” if V_+ and V_- can be bounded by some function of Z , which is usually a linear function. Functions of independent random variables which satisfy the self bounding property appear in reasonable amounts in the literature; examples include suprema of nonnegative empirical processes and conditional Rademacher averages. Further details and references about concentration inequalities for self-bounded functions of independent random variables are available in [18, 20] and [92].

We shall say a function f is self-bounded (w.r.t. X) if $v(X) \leq Bf(X) + C$ for some constants B and C , where v is defined in (3.1).

In Section 3.7, we shall provide some applications of this result to obtain Bernstein type concentration for functionals related to random permutations (matching problem and Spearman’s footrule).

Theorem 3.9 *Continuing with the notation introduced at the beginning of the Chapter, suppose B, C are finite positive constants such that $v(x) \leq Bf(x) + C$ for each x . Then $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-t^2/(2C+2Bt)}$ for any $t \geq 0$.*

Example. As usual, consider $X = \sum_{i=1}^n Y_i$, where Y_i ’s are now independent random variables taking value in $[0, 1]$. Let $\mu_i = \mathbb{E}(Y_i)$. We shall use Theorem 3.9 to prove a version of Bernstein’s inequality in this setup. As before, let $f(X) = X - \mathbb{E}(X)$. Let X' be constructed as in the example following Theorem 3.2. From previous computation, we know that

$$v(X) = \frac{1}{2} \sum_{i=1}^n \mathbb{E}((Y'_i - Y_i)^2 | X) = \frac{1}{2} \sum_{i=1}^n (\mathbb{E}Y_i^2 - 2\mu_i \mathbb{E}(Y_i | X) + \mathbb{E}(Y_i^2 | X)).$$

Using $Y_i^2 \leq Y_i$ (because $0 \leq Y_i \leq 1$), we get

$$v(X) \leq \frac{1}{2} \sum_{i=1}^n (\mathbb{E}(Y_i) + \mathbb{E}(Y_i | X)) = \frac{1}{2} (\mathbb{E}(X) + X).$$

Thus, taking $B = 1/2$ and $C = \mathbb{E}(X)$, we get

$$\mathbb{P}\{|X - \mathbb{E}(X)| \geq t\} \leq 2e^{-t^2/(2\mathbb{E}(X)+t)}.$$

Note, for instance, if $\mu_i = 1/2$ for all i , then $\mathbb{E}(X) = n/2$, and this bound is essentially equivalent to the Hoeffding bound. The increase in efficiency is apparent only when μ_i 's are small.

Proof of Theorem 3.9. Proceeding exactly as in the proof of Theorem 3.3, we get

$$\begin{aligned} |m'(\theta)| &\leq |\theta| \mathbb{E}(e^{\theta f(X)} v(X)) \\ &\leq |\theta| \mathbb{E}(e^{\theta f(X)} (Bf(X) + C)) \\ &= B|\theta| m'(\theta) + C|\theta| m(\theta). \end{aligned}$$

Since m is a convex function and $m'(0) = \mathbb{E}(f(X)) = 0$, therefore $m'(\theta)$ always has the same sign as θ . Thus, for $0 \leq \theta < 1/B$, the above inequality translates into

$$\frac{d}{d\theta} \log m(\theta) \leq \frac{C\theta}{1 - B\theta}.$$

Using this and recalling that $m(0) = 1$, we have

$$\log m(\theta) \leq \int_0^\theta \frac{Cu}{1 - Bu} du \leq \frac{C\theta^2}{2(1 - B\theta)}.$$

Putting $\theta = t/(C + Bt)$, we get

$$\mathbb{P}\{f(X) \geq t\} \leq \exp(-\theta t + \log m(\theta)) \leq e^{-t^2/(2C+2Bt)}.$$

The lower tail can be done similarly (though the argument is not exactly symmetric in this case). □

3.7 Example: Matching problem and Spearman's footrule

Suppose π is chosen uniformly at random from the set of all permutations of $\{1, \dots, n\}$. It is a classical probabilistic fact (the matching problem) that as $n \rightarrow \infty$, the distribution of the number of fixed points of π converges weakly to the Poisson distribution with mean 1. Error bounds can also be obtained by various methods. The question of tail bounds for this random variable, with explicit constants, is a reasonable problem to look at. Although this question can presumably be tackled by Talagrand's technique ([107], section 5; also discussed in section 2.6 here) for random permutations, it is a good test case for our theory.

Another interesting problem related to random permutations is the behavior of $\sum_{i=1}^n |i - \pi(i)|$. This statistic, known as Spearman's footrule (see, e.g. [57]), arises in the nonparametric theory of statistics.

In the rest of this section, we shall work out concentration inequalities for a generalized version of these statistics, originating in an early work of Hoeffding [51].

Let $\{a_{ij}\}$ be an $n \times n$ array of real numbers, assumed to be in $[0, 1]$ for our purposes. Let π be a random (uniform) permutation of $\{1, \dots, n\}$, and let $X = \sum_{i=1}^n a_{i\pi(i)}$. This class of random variables was first studied by Hoeffding [51], who proved that they are approximately normally distributed under certain conditions. The following proposition gives concentration inequalities in this setup:

Proposition 3.10 *Let $\{a_{ij}\}_{1 \leq i, j \leq n}$ be a collection of numbers from $[0, 1]$. Let $X = \sum_{i=1}^n a_{i\pi(i)}$, where π is drawn from the uniform distribution over the set of all permutations of $\{1, \dots, n\}$. Then for any $t \geq 0$, $\mathbb{P}\{|X - \mathbb{E}(X)| \geq t\} \leq 2e^{-t^2/(4\mathbb{E}(X)+2t)}$.*

Remarks. Note that the bound does not explicitly depend on n . This is a consequence of the assumption that the a_{ij} 's are bounded. A version of this theorem sans the boundedness assumption can also be proved using our techniques, but then the bound will involve n . The classical concentration result of Maurey [70], stated as Theorem 2.14 in the review section 2.6 cannot give a Bernstein type bound like the above. Talagrand's theorem ([107], Theorem 5.1) might, but it is not clear from the

abstract form whether it really does. McDiarmid's corollary [72] of Talagrand's result certainly cannot give a result like the above.

Example 1. (Matching problem.) Taking $a_{ij} = \mathbb{I}\{i = j\}$, we get X to be the number of fixed points of π . Since $\mathbb{E}(X) = 1$, we get the exponential tail bound $\mathbb{P}\{|X - 1| \geq t\} \leq 2e^{-t^2/(4+2t)}$, which does not depend on n .

Example 2. (Spearman's footrule.) Sometimes, the boundedness of the a_{ij} 's can be overcome by just dividing by the maximum. We give one such example now.

In nonparametric statistics, a standard measure of distance between two permutations π and σ is the Spearman's footrule, defined as

$$\rho(\pi, \sigma) := \sum_{i=1}^n |\pi(i) - \sigma(i)|.$$

A standard reference for the uses of Spearman's footrule in nonparametric statistics is the book [57] by Kendall and Gibbons. From a statistical point of view, it is of interest to know the distribution of the Spearman's footrule distance between the identity and a random uniform permutation. Diaconis and Graham [29] proved the following theorem:

Theorem 3.11 [Diaconis and Graham [29]] *Let $\rho(\pi, \sigma) = \sum_{i=1}^n |\pi(i) - \sigma(i)|$. If π is chosen uniformly, then*

$$\begin{aligned} \mathbb{E}(\rho) &= \frac{1}{3}(n^2 - 1), \\ \text{Var}(\rho) &= \frac{1}{45}(n+1)(2n^2 + 7), \quad \text{and} \\ \mathbb{P}\left\{\frac{\rho - \mathbb{E}(\rho)}{\sqrt{\text{Var}(\rho)}} \leq t\right\} &\rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-x^2/2} dx \end{aligned}$$

as $n \rightarrow \infty$, for each $t \in \mathbb{R}$.

We shall use Proposition 3.10 to get finite sample tail bounds of the correct order. Without loss of generality, we can take σ to be the identity permutation. Let

$$a_{ij} = \frac{|i - j|}{n}, \quad 1 \leq i, j \leq n.$$

Then $0 \leq a_{ij} \leq 1$. Let $X = \sum_{i=1}^n a_{i\pi(i)} = \rho/n$. Then by Proposition 3.10, $\mathbb{P}\{|X - \mathbb{E}(X)| \geq u\} \leq e^{-u^2/(4\mathbb{E}(X)+2u)}$. Now put $u = t\sqrt{\text{Var}(X)}$, and observe that

$$\frac{X - \mathbb{E}(X)}{\sqrt{\text{Var}(X)}} = \frac{\rho - \mathbb{E}(\rho)}{\sqrt{\text{Var}(\rho)}}.$$

By the Diaconis-Graham computation, $\mathbb{E}(X) = (n^2 - 1)/3n$ and $\text{Var}(X) = (n+1)(2 + 7n^{-2})/45$. Combining, we get the following tail bound:

Proposition 3.12 *Let ρ be the Spearman's footrule distance between the identity and a uniformly chosen permutation of $\{1, \dots, n\}$. For any $t \geq 0$, we have*

$$\mathbb{P}\left\{\left|\frac{\rho - \mathbb{E}(\rho)}{\sqrt{\text{Var}\rho}}\right| \geq t\right\} \leq 2e^{-t^2/(\alpha_n + \beta_n t)},$$

where

$$\begin{aligned} \alpha_n &= \frac{60(n-1)}{n(2+7n^{-2})} \rightarrow 30 \quad \text{as } n \rightarrow \infty, \quad \text{and} \\ \beta_n &= 2\sqrt{45}(n+1)^{-1/2}(2+7n^{-2})^{-1/2} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Note that our tail bound is approximately a gaussian bound (and hence, of the correct order) for large n . However, the constants are poor. The reason is that Proposition 3.10 is designed for random variables that exhibit Poissonian behavior. Our X in this problem does not belong to that class, particularly because its variance is smaller than its mean by a factor significantly smaller than unity. Thus, although Proposition 3.10 can be successfully applied to this problem to get concentration bounds of the correct order, it is not an ideal example.

Proof of Proposition 3.10. Construct X' as follows: Choose I, J uniformly and independently at random from $\{1, \dots, n\}$. Let $\pi' = \pi \circ (I, J)$, where (I, J) denotes the transposition of I and J . It can be easily verified that (π, π') is an exchangeable pair. Hence if we let

$$X' := \sum_{i=1}^n a_{i\pi'(i)},$$

then (X, X') is also an exchangeable pair. Now note that

$$\begin{aligned} \frac{1}{2}\mathbb{E}(n(X - X')|\pi) &= \frac{n}{2}\mathbb{E}(a_{I\pi(I)} + a_{J\pi(J)} - a_{I\pi(J)} - a_{J\pi(I)}|\pi) \\ &= \frac{1}{n} \sum_{i,j} a_{i\pi(i)} - \frac{1}{n} \sum_{i,j} a_{i\pi(j)} \\ &= X - \mathbb{E}(X). \end{aligned}$$

Thus, we can take $f(x) = x - \mathbb{E}(X)$ and $F(x, y) = \frac{1}{2}n(x - y)$. Now note that since $0 \leq a_{ij} \leq 1$ for all i and j , we have

$$\begin{aligned} v(X) &= \frac{n}{4}\mathbb{E}((X - X')^2|\pi) \\ &= \frac{1}{4n} \sum_{i,j} (a_{i\pi(i)} + a_{j\pi(j)} - a_{i\pi(j)} - a_{j\pi(i)})^2 \\ &\leq \frac{1}{2n} \sum_{i,j} (a_{i\pi(i)} + a_{j\pi(j)} + a_{i\pi(j)} + a_{j\pi(i)}) \\ &= X + \mathbb{E}(X) = f(X) + 2\mathbb{E}(X). \end{aligned}$$

Applying Theorem 3.9 with $B = 1$ and $C = 2\mathbb{E}(X)$ completes the proof. $\square$

3.8 Inequalities for unbounded differences

In this section, we present a some results which are applicable when $v(X)$ is unbounded. As usual, we shall supplement with trivial examples. A nontrivial application will be worked out in section 4.5 of the next chapter.

The first result of this section is an extension of Theorem 3.3 which only requires reasonable bounds on the moment generating function of $v(X)$, assuming that it exists. To be more specific, note that if $v(X)$ is bounded, then

$$\lim_{L \rightarrow \infty} L^{-1} \log \mathbb{E}(e^{Lv(X)}) = \|v(X)\|_{\infty}.$$

Thus, the number $r(L)$ defined as

$$r(L) := L^{-1} \log \mathbb{E}(e^{Lv(X)})$$

may serve as a surrogate for an actual bound on $v(X)$, in situations where such a bound does not exist, or is not representative of the true size of $v(X)$. The following theorem allows us the flexibility of using $r(L)$ with appropriate choice of L :

Theorem 3.13 *Suppose $r(L)$ is defined as above. Fix any $L > 0$. Then we have $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-t^2/(2r(L)+4tL^{-1/2})}$ for any $t \geq 0$.*

Remarks. The idea is to choose L so that $L^{-1/2} \ll r(L)$. In particular, observe that if $v(X) \leq C$ almost surely for some constant C , we can take $L \rightarrow \infty$ and get the bound in Theorem 3.3.

Example. Again, let $X = \sum_{i=1}^n Y_i$, where Y_i 's are independent zero mean random variables. However, we shall now drop the boundedness assumption, and assume only that Y_i 's have gaussian tails: In other words, assume that there exists $\theta > 0$ such that $\mathbb{E}(e^{\theta Y_i^2}) \leq K_\theta < \infty$ for each i for some fixed constant K_θ . Choosing $L = 2\theta$ and applying Jensen's inequality, we have

$$\begin{aligned} r(L) &\leq (2\theta)^{-1} \log \mathbb{E}(e^{\theta \sum_{i=1}^n (\mathbb{E}(Y_i^2) + Y_i^2)}) \\ &\leq (2\theta)^{-1} \sum_{i=1}^n 2 \log \mathbb{E}(e^{\theta Y_i^2}) \leq \theta^{-1} n \log K_\theta. \end{aligned}$$

The above result now gives the bound

$$\mathbb{P}\{|X| \geq t\} \leq 2 \exp\left(-\frac{t^2}{2\theta^{-1}n \log K_\theta + 4t(2\theta)^{-1/2}}\right).$$

Note how this reduces to the Hoeffding bound when the Y_i 's are bounded.

Proof of Theorem 3.13. Let $u(X) = e^{\theta f(X)}/m(\theta)$, where $m(\theta) = \mathbb{E}(e^{\theta f(X)})$, as

usual. Applying Jensen's inequality, we have for $\theta \geq 0$

$$\begin{aligned} m'(\theta) &\leq \theta \mathbb{E}(e^{\theta f(X)} v(X)) \\ &= L^{-1} \theta m(\theta) \mathbb{E} \left[u(X) \left(\log \frac{e^{Lv(X)}}{u(X)} + \log u(X) \right) \right] \\ &\leq L^{-1} \theta m(\theta) \log \mathbb{E}(e^{Lv(X)}) + L^{-1} \theta \mathbb{E}(e^{\theta f(X)} \log u(X)). \end{aligned}$$

Recall that $m'(0) = \mathbb{E}(f(X)) = 0$, $m(0) = 1$, and m is a convex function. Hence $m(\theta) \geq 1$ for all θ . Consequently, $\log u(X) \leq \theta f(X)$. Using this in the above bound, we have

$$m'(\theta) \leq r(L) \theta m(\theta) + L^{-1} \theta^2 m'(\theta).$$

Written differently, this gives

$$\frac{d}{d\theta} \log m(\theta) \leq \frac{r(L) \theta}{1 - L^{-1} \theta^2}$$

for $0 \leq \theta < L^{1/2}$. Now take any $t \geq 0$, and let θ be the positive root of the equation

$$\frac{r(L) \theta}{1 - L^{-1} \theta^2} = t.$$

Explicitly, we have

$$\theta = \frac{Lr(L)}{2t} \left(\sqrt{1 + \frac{4t^2}{r(L)^2 L}} - 1 \right). \quad (3.19)$$

Using the inequality $\sqrt{1 + \alpha} \leq 1 + \sqrt{\alpha}$, it is easy to see that $\theta < L^{1/2}$. Another application of the same inequality gives

$$\begin{aligned} \sqrt{1 + \alpha} - 1 - \frac{\alpha}{2(1 + \sqrt{\alpha})} &\geq \sqrt{1 + \alpha} - 1 - \frac{\alpha}{2\sqrt{1 + \alpha}} \\ &= \frac{(\sqrt{1 + \alpha} - 1)^2}{2\sqrt{1 + \alpha}} \geq 0. \end{aligned}$$

Using this inequality in (3.19), we get

$$\theta \geq \frac{t}{r(L) + 2tL^{-1/2}}.$$

Now, with the θ defined in (3.19) we have

$$\log m(\theta) \leq \int_0^\theta \frac{r(L)u}{1 - L^{-1}u^2} du \leq \frac{r(L)\theta^2}{2(1 - L^{-1}\theta^2)} = \frac{\theta t}{2}.$$

Thus, by Chebychev's inequality we get

$$\mathbb{P}\{f(X) \geq t\} \leq e^{-\theta t + \log m(\theta)} \leq e^{-\theta t/2}.$$

Using the lower bound on θ derived above, we get the desired expression. The lower tail bounds are obtained by symmetry. $\square$

Our next result is the exchangeable pairs version of the Burkholder-Davis-Gundy inequality [21, 22, 23] from classical probability. In its simplest form, this inequality says that for a martingale difference sequence $\{X_i\}_{1 \leq i \leq n}$ adapted to some filtration $\{\mathcal{F}_i\}_{1 \leq i \leq n}$, we have for any $p \geq 1$ the inequality

$$\mathbb{E} \left| \sum_{i=1}^n X_i \right|^{2p} \leq (2p-1)^p \mathbb{E} \left| \sum_{i=1}^n \mathbb{E}(X_i^2 | \mathcal{F}_{i-1}) \right|^p.$$

Note that the inequality is trivially an equality for $p = 1$. Essentially, this inequality gives a p^{th} moment expression of the notion that a martingale which is the sum of a homogeneous sequence of differences grows like $n^{1/2}$. (Analogously, the Hoeffding inequality gives a tail estimate expression of this notion when the differences are bounded.) This is clear from the observation that the right hand side above is bounded by $n^{p-1} \sum_{i=1}^n \mathbb{E}|X_i|^{2p} = O(n^p)$ by Jensen's inequality, and so if the X_i 's are of comparable sizes then $\mathbb{E}|\sum X_i|^{2p} = O(n^p)$, which "shows" that $\sum X_i = O(n^{1/2})$ upto p^{th} moment accuracy.

Recently, Boucheron, Bousquet, Lugosi & Massart [20] have derived a useful version of this inequality for general functions of independent random variables (rather than just sums). Moment bounds are often useful when we can take p to be large (growing with n), because the resulting Chebychev inequalities give surprisingly efficient tail bounds. In fact, it is an easy fact that for any suitable random variable X and any $t \geq 0$,

$$\mathbb{P}\{|X| \geq t\} \leq \inf_{p \geq 1} \frac{\mathbb{E}|X|^p}{t^p} \leq \inf_{\theta \geq 0} \frac{\mathbb{E}(e^{\theta|X|})}{e^{\theta t}},$$

which shows that optimized Chebychev bounds are better than optimized Chernoff bounds. For applications of the moment inequalities to obtain efficient tail bounds for a variety of complicated functions of independent random variables, we refer to [20].

We shall now present the exchangeable pairs version of the Burkholder-Davis-Gundy inequality. In the following, $\|\cdot\|_p$ will denote the L^p norm of random variables; that is, for a random variables Y , $\|Y\|_p := (\mathbb{E}|Y|^p)^{1/p}$.

Theorem 3.14 *For any positive integer p , we have*

$$\|f(X)\|_{2p}^2 \leq (2p-1)\|v(X)\|_p.$$

Example. To see why this can be called the Burkholder-Davis-Gundy inequality for exchangeable pairs, consider, as usual, sums of independent random variables. Let $Y_1, \dots, Y_n$ be independent mean zero random variables with finite p^{th} moment, where p is a fixed positive integer. Let $X = \sum_{i=1}^n Y_i$. Construct X' as in the previous sections, and recall that $f(X) = X$, $F(X, X') = n(X - X')$, and

$$v(X) = \frac{1}{2} \sum_{i=1}^n (\mathbb{E}(Y_i^2) + \mathbb{E}(Y_i^2|X)).$$

By the above theorem and simple applications of Minkowski and Jensen inequalities, we get

$$\|X\|_{2p}^2 \leq (2p-1) \left\| \sum_{i=1}^n Y_i^2 \right\|_p,$$

which is the classical inequality for sums of independent random variables.

Proof of Theorem 3.14. By Lemma 3.1, we have

$$\mathbb{E}(f(X)^{2p}) = \frac{1}{2}\mathbb{E}((f(X)^{2p-1} - f(X')^{2p-1})F(X, X')).$$

By the inequality

$$|x^{2p-1} - y^{2p-1}| \leq \frac{2p-1}{2}(x^{2p-2} + y^{2p-2})|x - y|$$

which follows easily from a convexity argument very similar to (3.3), we have

$$\mathbb{E}(f(X)^{2p}) \leq (2p-1)\mathbb{E}(f(X)^{2p-2}v(X))$$

By Hölder's inequality, we get

$$\mathbb{E}(f(X)^{2p}) \leq (2p-1)(\mathbb{E}(f(X)^{2p}))^{(p-1)/p}(\mathbb{E}(v(X)^p))^{1/p}.$$

The proof is completed by transferring $\mathbb{E}(f(X)^{2p})^{(p-1)/p}$ to the other side. $\square$

3.9 A refinement of the exponential inequality

The following is a refinement of Theorem 3.3 which allows us, in particular, to derive concentration bounds in the Sherrington-Kirkpatrick model of spin glasses in the next section.

Theorem 3.15 *Define*

$$\begin{aligned} v_1(x) &:= \frac{1}{2}\mathbb{E}((f(X) - f(X'))F(X, X')|X = x) \quad \text{and} \\ v_2(x) &:= \frac{1}{4}\mathbb{E}((f(X) - f(X'))^2|F(X, X')||X = x). \end{aligned}$$

Suppose C and ε are constants such that $|v_1(X)| \leq C$ and $v_2(X) \leq \varepsilon$ a.s. Then $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-Ct^2/(2C^2+8\varepsilon t)}$ for any $t \geq 0$.

Remark. Note the difference between v and v_1 : the latter has no absolute value inside. This can be considered as a “second order refinement” of the original inequality in Theorem 3.3.

Proof. Recall the identity

$$m'(\theta) = \frac{1}{2} \mathbb{E}((e^{\theta f(X)} - e^{\theta f(X')})F(X, X'))$$

that was derived using Lemma 3.1 in the proof of Theorem 3.3. Instead of simply using the inequality $|e^x - e^y| \leq \frac{1}{2}(e^x + e^y)|x - y|$, we now take a finer recourse: For any $x \leq y$, note that

$$\begin{aligned} 0 &\leq \frac{1}{2}(e^y + e^x)(y - x) - (e^y - e^x) \\ &= \int_x^y \left(\frac{e^y - e^x}{y - x}(u - x) + e^x - e^u \right) du \\ &\leq \int_x^y \left(\frac{1}{2}(e^y + e^x)(u - x) + 0 \right) du \\ &= \frac{1}{4}(e^y + e^x)(y - x)^2. \end{aligned}$$

Thus, we have

$$e^x - e^y = \frac{1}{2}(e^x + e^y)(x - y) + \delta(x, y),$$

where

$$|\delta(x, y)| \leq \frac{1}{4}(e^x + e^y)(x - y)^2.$$

Combining this with the observation that $(x - y)F(x, y) = (y - x)F(y, x)$, is it not difficult to deduce that for any $\theta \geq 0$ we have

$$\begin{aligned} m'(\theta) &\leq \theta \mathbb{E}(e^{\theta f(X)} v_1(X)) + \theta^2 \mathbb{E}(e^{\theta f(X)} v_2(X)) \\ &\leq (C\theta + \varepsilon\theta^2)m(\theta). \end{aligned}$$

This gives, for any $\theta \geq 0$,

$$\log m(\theta) \leq \frac{C\theta^2}{2} + \frac{\varepsilon\theta^3}{3}.$$

Now fix any $t \geq 0$. Let θ be the positive solution of the equation $\varepsilon\theta^2 + C\theta = t$. Explicitly,

$$\theta = \frac{C}{2\varepsilon} \left(\sqrt{1 + \frac{4\varepsilon t}{C^2}} - 1 \right).$$

Then

$$\begin{aligned} -\theta t + \log m(\theta) &\leq -\theta(\varepsilon\theta^2 + C\theta) + \frac{C\theta^2}{2} + \frac{\varepsilon\theta^3}{3} \\ &= -\frac{C\theta^2}{2} - \frac{2\varepsilon\theta^3}{3} \leq -\frac{C\theta^2}{2}. \end{aligned}$$

Now note that for any $\alpha \geq 0$,

$$\sqrt{1 + \alpha} - 1 - \frac{\alpha}{2\sqrt{1 + \alpha}} = \frac{(\sqrt{1 + \alpha} - 1)^2}{2\sqrt{1 + \alpha}} \geq 0.$$

Thus,

$$\theta \geq \frac{C}{2\varepsilon} \left(\frac{\frac{4\varepsilon t}{C^2}}{2\sqrt{1 + \frac{4\varepsilon t}{C^2}}} \right) = \frac{t}{\sqrt{C^2 + 4\varepsilon t}}.$$

Combining the steps, we get

$$\mathbb{P}\{f(X) \geq t\} \leq e^{-\theta t + \log m(\theta)} \leq e^{-C\theta^2/2} \leq e^{-Ct^2/(2C^2 + 8\varepsilon t)}.$$

The lower tail bound follows by symmetry. □

3.10 Application to spin glasses

The Sherrington-Kirkpatrick (S-K) model of spin glasses considers the Hamiltonian

$$H(\sigma) = -n^{-1/2} \sum_{1 \leq i < j \leq n} g_{ij} \sigma_i \sigma_j - h \sum_{i=1}^n \sigma_i,$$

where g_{ij} 's are a fixed realization of a collection of i.i.d. standard gaussian random variables. For recent advances in the rigorous analysis of this model, we refer to Chapter 2 of the book [111] by Michel Talagrand, and also his proof [112] of the

Parisi formula for the limiting free energy in this model. For the physical perspective, one should look at the book [74] by Mézard, Parisi and Virasoro.

In spite of the fact that the Parisi formula has been proved, very little is explicitly known about the low temperature phase of the S-K model at the time of writing this thesis. In fact, as Talagrand admits ([111], p. 182), “we do not even know where to start”!

The mean field equations discussed in section 3.4 are no longer valid in the S-K model, even at high temperatures. Instead, physicists think that a modification of the naive mean field equations (3.6), the so-called TAP equations, hold for the S-K model. These are as follows:

$$\langle \sigma_i \rangle = \tanh \left(\frac{\beta}{\sqrt{n}} \sum_{j \leq n, j \neq i} g_{ij} \langle \sigma_j \rangle + h - \beta^2(1-q) \langle \sigma_i \rangle \right), \quad i = 1, \dots, n \quad (3.20)$$

where q solves $q = \mathbb{E} \tanh^2(\beta Z \sqrt{q} + h)$, Z being a standard gaussian random variable. The validity of this conjecture has been established rigorously for sufficiently small β by Talagrand ([111], Theorem 2.4.20).

In this section, we shall prove that another set of mean field type equations are valid, irrespective of the temperature, in a class of models which will encompass the S-K model. For instance, we shall be able to show that in this class of models, the magnetization $m(\sigma) = \frac{1}{n} \sum_{i=1}^n \sigma_i$ satisfies

$$m(\sigma) \approx \frac{1}{n} \sum_{i=1}^n \tanh(\beta m_i(\sigma)) \quad \text{with high probability,} \quad (3.21)$$

where $m_i(\sigma) := n^{-1/2} \sum_{j \leq n, j \neq i} g_{ij} \sigma_j + h$ is the “local field” at site i . It will follow from the same argument that whenever $k \ll \sqrt{n}$, we also have

$$\frac{1}{n} \sum_{i=1}^n \prod_{r=1}^k \sigma_i^r \approx \frac{1}{n} \sum_{i=1}^n \prod_{r=1}^k \tanh(\beta m_i(\sigma^r)), \quad (3.22)$$

where $\sigma^1, \dots, \sigma^k$ are i.i.d. from the Gibbs measure. The left hand side, called the “overlap” of the k configurations, is a particularly interesting quantity in the S-K

model when $k = 2$.

However, we must admit that we have found no use for all this information till now. Still, it may be interesting just because the low temperature phase of the S-K model is a highly intractable object.

Our results will be valid whenever the L^2 operator norms of the matrices J and $J_2 := (J_{ij}^2)$ are bounded. Recall that the L^2 operator norm of a matrix A is defined as

$$\|A\| := \max_{\|x\|=1} \|Ax\|.$$

Alternatively, it is the square root of the maximum eigenvalue of $A^T A$. In the S-K model, it is well-known that the norm of J is bounded in probability. In fact, it is known that with $J^{(n)} = (n^{-1/2}g_{ij})_{1 \leq i,j \leq n}$, where $\{g_{ij}\}_{1 \leq i \leq j \leq n}$ are i.i.d. standard gaussian random variables and $g_{ji} = g_{ij}$, we have

$$\|J^{(n)}\| \rightarrow 2 \text{ in probability.}$$

A proof of this result, as well as tail bounds, can be found in the survey article by Bai [8], section 2.2. It is also easy to see that the norm of J_2 is bounded in probability in the S-K model, because

$$\|J_2\| \leq \left[n \max_{1 \leq i \leq n} \sum_{j=1}^n J_{ij}^4 \right]^{1/2} = \left[\max_{1 \leq i \leq n} \sum_{j=1}^n \frac{g_{ij}^4}{n} \right]^{1/2} \rightarrow \mathbb{E}(g_{11}^4) \text{ in probability,}$$

since for each i , $\frac{1}{n} \sum_{j=1}^n g_{ij}^4$ is concentrated around $\mathbb{E}(g_{11}^4)$ and the tails fall off sharply enough for the above to hold (a fact that can be easily proved using the Burkholder-Davis-Gundy inequality). The key result of this section is the following:

Theorem 3.16 *Let J be the interaction matrix in a model described by the Hamiltonian (3.4). Let $J_2 = (J_{ij}^2)_{1 \leq i,j \leq n}$. Take any $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$. Fix $\beta \geq 0$, and let*

$$C = C(\alpha, \beta) := 2(1 + \beta\|J\| + \beta^2\|J_2\|)\|\alpha\|^2, \quad \text{and} \\ \varepsilon = \varepsilon(\alpha, \beta) := 4\left(\max_{1 \leq i \leq n} |\alpha_i|\right) [1 + (\beta\|J\| + \beta^2\|J_2\|)^2] \|\alpha\|^2.$$

For each i , let $m_i = m_i(\sigma) := \sum_{j=1}^n J_{ij}\sigma_j + h_i$. Let

$$Y = \sum_{i=1}^n \alpha_i (\sigma_i - \tanh(\beta m_i)).$$

If σ is drawn from the Gibbs measure at inverse temperature β , then $\mathbb{E}(Y) = 0$, $\text{Var}(Y) \leq C$, and for any $t \geq 0$, $\mathbb{P}\{|Y| \geq t\} \leq 2e^{-Ct^2/(2C^2+8\epsilon t)}$.

Applications. Taking $\alpha_i = 1/n$ for each i , we get $\|\alpha\|^2 = 1/n$. Fix $\beta \geq 0$, and let $K = \beta\|J\| + \beta^2\|J_2\|$. As observed before, $K = O(1)$ in the S-K model. We have $C = (2 + 2K)/n$ and $\epsilon = 4(1 + K^2)/n^2$. Thus, if we let $m(\sigma) := \frac{1}{n} \sum_{i=1}^n \sigma_i$ be the magnetization of a configuration drawn from the Gibbs measure at inverse temperature β , and let

$$Y = m(\sigma) - \frac{1}{n} \sum_{i=1}^n \tanh(\beta m_i(\sigma)),$$

where, as usual, $m_i(\sigma) = \sum_{j=1}^n J_{ij}\sigma_j + h_i$ is the local field at i , then $\mathbb{E}(Y^2) \leq (2 + 2K)/n$ and

$$\mathbb{P}\{|Y| \geq t\} \leq 2e^{-nt^2/(a+bt)}, \quad (3.23)$$

where $a = 4 + 4K$ and $b = 16(1 + K^2)/(1 + K)$ are free of n . This shows (3.21).

The relation (3.22) for the k^{th} -order overlaps can be treated exactly in the same way, by successively conditioning on $(\sigma^1, \dots, \sigma^{r-1}, \sigma^{r+1}, \dots, \sigma^k)$, $r = 1, \dots, k$, and getting

$$\frac{1}{n} \sum_{i=1}^n \prod_{s=1}^{r-1} \tanh(\beta m_i(\sigma^{s-1})) \prod_{s=r}^k \sigma_i^s \approx \frac{1}{n} \sum_{i=1}^n \prod_{s=1}^r \tanh(\beta m_i(\sigma^{s-1})) \prod_{s=r+1}^k \sigma_i^s$$

at the r^{th} stage. Roughly, this can be carried out till $k = o(\sqrt{n})$, since the errors accrued at each stage are like $n^{-1/2}$.

Proof of Theorem 3.16. As in the proof of Proposition 3.4, we construct an exchangeable pair by taking a step in the Gibbs sampler chain as follows: First, draw

σ from the Gibbs measure. Next, choose a coordinate I uniformly at random, and replace the I^{th} coordinate of σ by a sample drawn from the conditional distribution of the σ_I given $\{\sigma_j, j \neq I\}$. Let

$$F(\sigma, \sigma') := n \sum_{i=1}^n \alpha_i (\sigma_i - \sigma'_i) = n(\sigma_I - \sigma'_I).$$

Then F is antisymmetric and

$$\begin{aligned} f(\sigma) &= \mathbb{E}(F(\sigma, \sigma') | \sigma) = \sum_{i=1}^n \alpha_i (\sigma_i - \mathbb{E}(\sigma_i | \{\sigma_j, j \neq i\})) \\ &= \sum_{i=1}^n \alpha_i (\sigma_i - \tanh(\beta m_i(\sigma))). \end{aligned}$$

For any $\sigma \in \{-1, 1\}^n$ and $1 \leq j \leq n$, let

$$\sigma^{(j)} := (\sigma_1, \dots, \sigma_{i-1}, -\sigma_i, \sigma_{i+1}, \dots, \sigma_n).$$

Define $h : \mathbb{R} \rightarrow \mathbb{R}$ as $h(x) := \tanh(\beta x)$, and for each i, j , let

$$b_{ij} = b_{ij}(\sigma) := h(m_i(\sigma)) - h(m_i(\sigma^{(j)})),$$

where $m_i(\sigma) = \sum_{j=1}^n J_{ij} \sigma_j + h_i$, as defined in the statement of the theorem. Now note that

$$f(\sigma) - f(\sigma^{(j)}) = 2\alpha_j \sigma_j - \sum_{i=1}^n \alpha_i b_{ij}, \quad \text{and} \quad F(\sigma, \sigma^{(j)}) = 2n\alpha_j \sigma_j. \quad (3.24)$$

Let $p_j = p_j(\sigma) = \mathbb{P}\{\sigma'_j = -\sigma_j \mid \sigma, I = j\}$. Combining everything, we have

$$v_1(\sigma) = \frac{1}{2} \mathbb{E}((f(\sigma) - f(\sigma')) F(\sigma, \sigma') | \sigma) = 2 \sum_{j=1}^n \alpha_j^2 p_j - \sum_{i,j} \alpha_i b_{ij} \alpha_j \sigma_j p_j. \quad (3.25)$$

It is easy to verify that $\|h''\|_\infty \leq \beta^2$. Therefore,

$$\begin{aligned} & |h(m_i(\sigma)) - h(m_i(\sigma^{(j)})) - (m_i(\sigma) - m_i(\sigma^{(j)}))h'(m_i(\sigma))| \\ & \leq \frac{\beta^2}{2}(m_i(\sigma) - m_i(\sigma^{(j)}))^2. \end{aligned}$$

Now let $c_i = c_i(\sigma) := h'(m_i(\sigma))$, and note that $m_i(\sigma) - m_i(\sigma^{(j)}) = 2J_{ij}\sigma_j$. Combining, we have

$$|b_{ij} - 2J_{ij}\sigma_j c_i| \leq 2\beta^2 J_{ij}^2. \quad (3.26)$$

Finally, note that $|c_i| \leq \beta$. Using all this information, we get, for any $x, y \in \mathbb{R}^n$,

$$\begin{aligned} \left| \sum_{i,j} x_i y_j b_{ij} \right| & \leq \left| \sum_{i,j} x_i y_j (2J_{ij}\sigma_j c_i) \right| + \left| \sum_{i,j} x_i y_j (b_{ij} - 2J_{ij}\sigma_j c_i) \right| \\ & \leq 2\|J\| \left(\sum_{i=1}^n (x_i c_i)^2 \right)^{1/2} \left(\sum_{j=1}^n (y_j \sigma_j)^2 \right)^{1/2} + \sum_{i,j} |x_i y_j| 2\beta^2 J_{ij}^2 \\ & \leq (2\beta\|J\| + 2\beta^2\|J_2\|)\|x\|\|y\|. \end{aligned} \quad (3.27)$$

Let $K = 2\beta\|J\| + 2\beta^2\|J_2\|$. Using the above inequality in equation (3.25) with $x_i = \alpha_i$ and $y_j = \alpha_j \sigma_j p_j$, we get

$$|v_1(\sigma)| \leq (2 + K) \sum_{i=1}^n \alpha_i^2.$$

The first assertion of this theorem now follows by Theorem 3.2. Next, we use (3.24) once again (and the inequality $(a + b)^2 \leq 2a^2 + 2b^2$) to get

$$\begin{aligned} v_2(\sigma) & = \frac{1}{4} \mathbb{E}((f(\sigma) - f(\sigma'))^2 | F(\sigma, \sigma') | | \sigma) \\ & \leq 4 \sum_{i=1}^n |\alpha_i|^3 + \sum_{j=1}^n |\alpha_j| \left(\sum_{i=1}^n \alpha_i b_{ij} \right)^2 \\ & \leq (\max_i |\alpha_i|) \left[4 \sum_{i=1}^n \alpha_i^2 + \sum_{j=1}^n \left(\sum_{i=1}^n \alpha_i b_{ij} \right)^2 \right]. \end{aligned} \quad (3.28)$$

Now, put $x_i = \alpha_i$ and $y_j = \sum_{k=1}^n \alpha_k b_{kj}$, $i, j = 1, \dots, n$. Then

$$\begin{aligned} \|y\|^2 &= \sum_{j=1}^n \left(\sum_{i=1}^n \alpha_i b_{ij} \right)^2 = \sum_{i,j} x_i y_j b_{ij} \\ &\leq K \|x\| \|y\| \quad \text{by inequality (3.27).} \end{aligned}$$

Thus, $\|y\| \leq K \|x\|$, and therefore by (3.28) we have

$$v_2(\sigma) \leq 2(\max_i |\alpha_i|)(1 + K^2) \sum_{i=1}^n \alpha_i^2.$$

The proof is now completed by applying Theorem 3.15. □

Chapter 4

Theory and examples: Part II

We begin this chapter with Stein's observation [102] that an exchangeable pair (X, X') automatically defines a reversible Markov kernel P as

$$Pf(x) := \mathbb{E}(f(X')|X = x), \quad (4.1)$$

where f is any function such that $\mathbb{E}|f(X)| < \infty$. All other notation will be the same as what was defined at the beginning of Chapter 3.

The main purpose of the subsequent discussion is to connect the concentration properties of the distribution of X with the rate of convergence to stationarity of a Markov chain following the kernel P . In particular, information about the rate of decay of $P^k f(x) - P^k f(y)$, where (x, y) is any point in the support of (X, X') , can give us a bound on $F(X, X')$ in a way that we are going to describe in the following pages. This is a direction that has not been systematically explored in the concentration literature, to the best of our knowledge. We shall also describe a coupling technique for getting a handle on the antisymmetric function F when f is known.

The techniques so developed, will be applied in section 4.2 to get concentration under a famous “weak dependence” condition from statistical physics. Examples from classical spin systems at high temperature and random proper graph colorings are worked out in the subsequent sections.

In section 4.5, we shall use the results of this section in another direction to obtain

concentration of Haar measures on compact groups based on convergence rates of random walks. The result will be used in section 4.6 to derive a quantitative version of Voiculescu's connection [114] between random matrices and free probability theory.

4.1 Explicit construction of $F(X, X')$

Our construction of F will involve the *Poisson equation* associated with the kernel P , which can be written as follows:

$$g - Pg = f. \tag{4.2}$$

Here f is a given function, and the objective is to solve for g . The Poisson equation is an object of deep mathematical significance. Its importance in the theory of Markov chains was realized after the work of Paul Meyer [73] and the subsequent investigation of Neveu [78]. The contributions by Nummelin [79] are also significant. A classical textbook reference is the book by Revuz [91]. Poisson's equation has been used in the probability literature by too many authors to mention in this short space; for a recent survey of the literature about Poisson's equation in the context of discrete Markov chains, we refer to Makowski and Shwartz [64].

The way we can think of using the solution to Poisson's equation in our problem is the following: If we let $F(x, y) = g(x) - g(y)$, where g is a solution to (4.2), then F is antisymmetric and

$$\mathbb{E}(F(X, X')|X) = g(X) - \mathbb{E}(g(X')|X) = g(X) - Pg(X) = f(X).$$

Generally, the solution to Poisson's equation is given by

$$g = \sum_{k=0}^{\infty} P^k f,$$

but problems with convergence are not uncommon. The solution need not be unique, either. Various conditions have been proposed for the existence and uniqueness of

solutions of Poisson's equation in various situations. For a summary of such results, we refer to [64].

We shall not need the weakest conditions for the existence of solutions to Poisson's equation in any of our applications. The following lemma formalizes our construction of F under generous assumptions, which are satisfied for geometrically ergodic Markov chains, and all of our examples belong to that class.

Lemma 4.1 *Let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a measurable function such that $\mathbb{E}f(X) = 0$. Suppose there is a finite constant L such that*

$$\sum_{k=0}^{\infty} |P^k f(x) - P^k f(y)| \leq L \quad \text{for every } x \text{ and } y. \quad (4.3)$$

Then the function

$$F(x, y) := \sum_{k=0}^{\infty} (P^k f(x) - P^k f(y))$$

satisfies $F(X, X') = -F(X', X)$ and $\mathbb{E}(F(X, X')|X) = f(X)$.

Proof. The convergence of the series defining F follows from the summability assumption (4.3). Also, F is clearly antisymmetric. Now,

$$\mathbb{E}(P^k f(X) - P^k f(X')|X) = P^k f(X) - P^{k+1} f(X).$$

Thus, for any N , we have

$$\sum_{k=0}^N \mathbb{E}(P^k f(X) - P^k f(X')|X) = f(X) - P^{N+1} f(X).$$

The condition (4.3) ensures that the above partial sums converge everywhere. Consequently, the sequence $\{P^{N+1} f(X)\}_{N \geq 0}$ also converges everywhere. Again, condition (4.3) implies that the limit is a constant, since for any x and y , $P^k f(x) - P^k f(y) \rightarrow 0$ as $k \rightarrow \infty$. Since $\mathbb{E}F(X, X') = 0 = \mathbb{E}f(X)$, this constant can only be zero. $\square$

Although Lemma 4.1 gives an explicit expression for F , it is somewhat inconvenient for practical purposes. We shall now give a coupling version of Lemma 4.1, that will be easier to work with in practice.

Let $\{X_k\}_{k \geq 0}$ and $\{X'_k\}_{k \geq 0}$ be two chains from the kernel defined by (X, X') , with arbitrary initial values, and coupled according to some coupling scheme which satisfies the following property:

P For every initial value (x, y) , and every k , the marginal distribution of X_k depends only on x and the marginal distribution of X'_k depends only on y .

Under this assumption, we have the following lemma:

Lemma 4.2 *Suppose the chains $\{X_k\}$ and $\{X'_k\}$ satisfy the property **P** described above. Let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a function such that $\mathbb{E}f(X) = 0$. Suppose there exists a finite constant L such that for every $(x, y) \in \mathcal{X}^2$,*

$$\sum_{k=0}^{\infty} |\mathbb{E}(f(X_k) - f(X'_k) | X_0 = x, X'_0 = y)| \leq L. \quad (4.4)$$

Then, the function F defined as

$$F(x, y) := \sum_{k=0}^{\infty} \mathbb{E}(f(X_k) - f(X'_k) | X_0 = x, X'_0 = y)$$

satisfies $F(X, X') = -F(X', X)$ and $\mathbb{E}(F(X, X') | X) = f(X)$.

Remark. In practice, we will start our chains with $X_0 = X$ and $X'_0 = X'$ for directly obtaining bounds on $F(X, X')$ in the process of verifying (4.4).

Proof. Property **P** implies that

$$\mathbb{E}(f(X_k) | X_0 = x, X'_0 = y) = \mathbb{E}(f(X_k) | X_0 = x) = P^k f(x),$$

where P is the kernel defined in (4.1). Similarly, $\mathbb{E}(f(X'_k) | X_0 = x, X'_0 = y) = P^k f(y)$.

The rest is a rewriting of the conclusions of Lemma 4.1. $\square$

Example. Let $\mathcal{X} = \mathbb{R}^n$, and let $X = (X_{(1)}, \dots, X_{(n)})$ be a vector with independent components. (The subscripts are put in brackets because we will be using $\{X_k\}_{k \geq 0}$ to denote a Markov chain on $\mathbb{R}^n$.) As a preliminary example, and also as a prelude to section 4.2, we shall now work out the concentration of $f(X)$, where f is a function satisfying

$$|f(x) - f(y)| \leq \sum_{i=1}^n c_i \mathbb{I}\{x_i \neq y_i\}.$$

In other words, if x and y differ only at coordinate i , then $|f(x) - f(y)| \leq c_i$. Our technique will give an easy way to recover a version of the bounded difference inequality (Theorem 2.3) without using martingale or information theoretic results. We shall only give a brief description of the steps in our solution, because the details will be worked out under a more general setting in the next section.

We produce X' by choosing a coordinate I uniformly at random, and replacing the I^{th} coordinate of X by $X_{(I)}^*$, where the vector $(X_{(1)}^*, \dots, X_{(n)}^*)$ has the same distribution as X and is independent of X . It is easy to see that (X, X') is an exchangeable pair.

The coupling is done in the natural way: At every step, choose the same coordinate I for both chains, and replace the I^{th} coordinate of both by the same realization of $X_{(I)}^*$. Since the number of coordinates at which X_k and X'_k differ cannot increase with k , it is clear (by the coupon collector's problem) that the coupling time for the chains has expected value bounded by $n \log n$. This proves condition (4.4).

Now suppose we start with $(X_0, X'_0) = (x, y)$. If x and y differ only at coordinate i , then $|f(X_k) - f(X'_k)| \leq c_i$ for every k , and the expected coupling time is n . Thus, by the representation of F in Lemma 4.2, we get $|F(x, y)| \leq nc_i$. Consequently, we have

$$\begin{aligned} v(X) &= \frac{1}{2} \mathbb{E}(|(f(X) - f(X'))F(X, X')||X) \\ &\leq \frac{1}{2n} \sum_{i=1}^n (c_i)(nc_i) = \frac{1}{2} \sum_{i=1}^n c_i^2. \end{aligned}$$

Theorem 3.3 now gives us a version of the well-known bounded difference inequality

(Theorem 2.3), albeit with a missing 2 in the exponent.

4.2 Concentration under weak dependence

Let $\mathcal{X} = \Omega^n$, where Ω is a Polish space. Consequently, our random variable X will now have n coordinates, which we shall assume to be *weakly dependent* in the sense of Dobrushin, a familiar notion from statistical mechanics which we shall define later in this section.

We shall also assume that our function f satisfies a Lipschitz condition with respect to a generalized Hamming distance on $\mathcal{X}$:

$$|f(x) - f(y)| \leq \sum_{i=1}^n c_i \mathbb{I}\{x_i \neq y_i\} \quad (4.5)$$

for some fixed constants $c_1, \dots, c_n$. As mentioned before, this just means that the value of the function does not change by more than c_i if the i^{th} coordinate is altered.

The generalization of the Hamming metric is useful because it allows us to get concentration for lower dimensional marginals. For instance, if we want the concentration of a function of $(X_1, \dots, X_k)$, where $k \leq n$, we can just consider every function of $(X_1, \dots, X_k)$ as a function of $(X_1, \dots, X_n)$ with $c_{k+1} = \dots = c_n = 0$.

In the following, we shall be using the notation $\bar{x}^i$ to denote the element of Ω^{n-1} obtained by omitting the i^{th} coordinate of the vector $x \in \Omega^n$.

We shall let μ denote the law of X , and for each i and x , $\mu_i(\cdot | \bar{x}^i)$ will stand for the law of the i^{th} coordinate of X given that $\bar{X}^i = \bar{x}^i$.

At this point, let us also mention that we shall usually denote the i^{th} coordinate of a vector x by x_i . Unfortunately, we are also using the notation X_k for the k^{th} element of a Markov chain. We hope that the context will clarify any ambiguity. In particular, to denote the i^{th} coordinate of X_k we shall use the notation $X_{k,i}$.

Finally, let us recall that for a square matrix A , the L^2 operator norm of A is defined as:

$$\|A\|_2 := \max_{\|y\|=1} \|Ay\|.$$

While we are at it, let us also recall that the L^∞ and L^1 operator norms can be expressed as

$$\|A\|_\infty := \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|, \quad \text{and} \quad \|A\|_1 := \max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}|.$$

It is a frequently useful fact that $\|A\|_2 \leq \sqrt{\|A\|_1 \|A\|_\infty}$.

The following theorem gives a simple way of obtaining concentration bounds for $f(X)$ under a contractivity condition on the conditional laws:

Theorem 4.3 *Suppose $A = (a_{ij})$ is an $n \times n$ matrix with nonnegative entries and zeros on the diagonal such that for any i , and any $x, y \in \Omega^n$,*

$$d_{TV}(\mu_i(\cdot | \bar{x}^i), \mu_i(\cdot | \bar{y}^i)) \leq \sum_{j=1}^n a_{ij} \mathbb{I}\{x_j \neq y_j\},$$

where d_{TV} is the total variation distance on the space of probability measures on Ω . Suppose f satisfies the generalized Lipschitz condition (4.5). If $\|A\|_2 < 1$, we have

$$\mathbb{P}\{|f(X) - \mathbb{E}f(X)| \geq t\} \leq 2 \exp\left(-\frac{(1 - \|A\|_2)t^2}{\sum_i c_i^2}\right)$$

for each $t \geq 0$.

Remarks. The matrix A is called “Dobrushin’s interdependence matrix” and the condition $\|A\|_\infty < 1$ is called Dobrushin’s condition (as introduced by Dobrushin [35] and extended by Dobrushin & Shlosman [36, 37]). Dobrushin’s condition ensures, among other things, the uniqueness of Gibbs states at high temperature. In a series of important papers, Stroock and Zegarlinski [103, 104, 105] showed that for spin systems on a lattice, the Dobrushin-Shlosman conditions are equivalent to the validity of logarithmic Sobolev inequalities for the associated Glauber dynamics. Though concentration inequalities follow from log-Sobolev inequalities, explicit constants are not available from this body of work. Our approach gives a direct way of getting explicit concentration bounds from Dobrushin type conditions. For more on Dobrushin’s condition and its consequences, see Georgii’s book [45], Chapter 8. For

a recent treatise on logarithmic Sobolev inequalities in the context of spin systems and Dobrushin-Shlosman type conditions, see Ledoux [62].

As mentioned in section 2.4, Dobrushin type conditions were recently used by Marton [67, 68] to obtain transportation cost inequalities (and hence concentration) under weak dependence. However, Marton's results do not seem to be suited for the setting that we are now working under. Besides, they require us to know some bounds on certain log-Sobolev constants, which are hard to obtain.

Proof of Theorem 4.3. To prove this theorem, we shall construct a reversible Markov kernel and a suitable coupling, and then directly apply Lemma 4.2 in conjunction with the tools from Chapter 3.

First, define a reversible Markov kernel as follows: At each step, choose a coordinate I uniformly from $\{1, \dots, n\}$. Then replace the I^{th} coordinate of the current state of the chain by an element of Ω chosen according to the conditional distribution of the I^{th} coordinate given the values of the other coordinates. This is the usual Gibbs sampling technique, and it is well-known and easy to prove that the chain is reversible. This is also known as the ‘‘Glauber dynamics’’ in the case of spin systems.

Now, we describe the coupling. Suppose at any stage, the X -chain is at x , and the X' -chain is at y . Choose a coordinate I uniformly at random. By the well-known property of the total variation distance, we can have two $\mathcal{X}$ -valued random variables W_1^I and W_2^I defined on some probability space such that $W_1^I \sim \mu_I(\cdot | \bar{x}^I)$, $W_2^I \sim \mu_I(\cdot | \bar{y}^I)$, and

$$\begin{aligned} \mathbb{P}\{W_1^I \neq W_2^I\} &= d_{TV}(\mu_I(\cdot | \bar{x}^I), \mu_I(\cdot | \bar{y}^I)) \\ &\leq \sum_{j=1}^n a_{Ij} \mathbb{I}\{\bar{x}_j^I \neq \bar{y}_j^I\} \quad (\text{by assumption}). \end{aligned}$$

Having obtained W_1^I and W_2^I , update the X -chain by putting W_1^I as the I^{th} coordinate (keeping all other coordinates the same) and update the X' -chain the same way using W_2^I . This is perhaps the most naive way to couple dependent variables, and is commonly known as ‘‘the greedy coupling’’.

Of course, people know how to make the above construction completely mathematically precise, and so we prefer to gloss over that aspect. But if someone is interested, he can look up the recent paper of Marton [68], where a more complex coupling has been handled in a very mathematically precise manner.

It is clear from the definition that this coupling indeed gives the Gibbs sampler chains as the marginals. To see that property **P** holds, note that for any i , the distribution of W_1^i as defined above depends only on the current state of the X -chain, irrespective of the status of the X' -chain. Similarly, the distribution of W_2^i depends only on the current state of the X' -chain. Thus, the distribution of X_{k+1} given (X_k, X'_k) depends only on X_k . Since the coupling is Markovian, we can proceed by induction to get a proof of property **P** for this coupling.

Now assume without loss of generality that $\mathbb{E}f(X) = 0$. The summability condition (4.4) will be verified later in the proof.

Fix $k \geq 0$, and suppose we used the above scheme to move to the $(k+1)^{\text{th}}$ stage. Then for any i ,

$$\mathbb{P}\{X_{k+1,i} \neq X'_{k+1,i} \text{ and } I \neq i | X_k, X'_k\} = \left(1 - \frac{1}{n}\right) \mathbb{I}\{X_{k,i} \neq X'_{k,i}\},$$

and

$$\begin{aligned} & \mathbb{P}\{X_{k+1,i} \neq X'_{k+1,i} \text{ and } I = i | X_k, X'_k\} \\ &= \frac{1}{n} \mathbb{P}\{W_1^i \neq W_2^i\} \leq \frac{1}{n} \sum_{j=1}^n a_{ij} \mathbb{I}\{X_{k,j} \neq X'_{k,j}\}. \end{aligned}$$

Adding up, and taking expectation given (X_0, X'_0) on both sides, we get

$$\begin{aligned} & \mathbb{P}\{X_{k+1,i} \neq X'_{k+1,i} | X_0, X'_0\} \\ & \leq \left(1 - \frac{1}{n}\right) \mathbb{P}\{X_{k,i} \neq X'_{k,i} | X_0, X'_0\} + \frac{1}{n} \sum_{j=1}^n a_{ij} \mathbb{P}\{X_{k,j} \neq X'_{k,j} | X_0, X'_0\}. \end{aligned}$$

For each k , let $\ell_k = \ell_k(X_0, X'_0)$ be the random vector whose i^{th} component is $\mathbb{P}\{X_{k,i} \neq X'_{k,i} | X_0, X'_0\}$. Let $B = (1 - \frac{1}{n})I + \frac{1}{n}A$, where I denotes the identity matrix. The above

inequality shows that $\ell_{k+1} \leq B\ell_k$ for every k , where ' $x \leq y$ ' means that $x_i \leq y_i$ for each coordinate i . Since everything is nonnegative, we can continue by induction to see that

$$\ell_k \leq B^k \ell_0.$$

Now let $c = (c_1, \dots, c_n)$ be the vector of constants from (4.5). Note that

$$\|B\|_2 \leq 1 - \frac{1}{n} + \frac{\|A\|_2}{n} < 1, \quad (4.6)$$

and hence, by the generalized Lipschitz property of f ,

$$\sum_{k=0}^{\infty} |\mathbb{E}(f(X_k) - f(X'_k) | X_0, X'_0)| \leq \sum_{k=0}^{\infty} \mathbb{E}(c \cdot \ell_k | X_0, X'_0) \leq \|c\| \|\ell_0\| \sum_{k=0}^{\infty} \|B\|^k < \infty.$$

This establishes condition (4.4), and so we can now invoke Lemma 4.2. Produce X' by starting from X and taking one step according to the Gibbs sampler kernel. Putting $X_0 = X$ and $X'_0 = X'$, we get

$$F(X, X') = \sum_{k=0}^{\infty} \mathbb{E}(f(X_k) - f(X'_k) | X_0, X'_0)$$

by Lemma 4.2. Thus, if v is defined as in (3.1), then

$$v(X) \leq \frac{1}{2} \sum_{k=0}^{\infty} \mathbb{E}(|(f(X_0) - f(X'_0))(f(X_k) - f(X'_k))| | X_0). \quad (4.7)$$

Now, by our definition of ℓ_k and the property (4.5) of f , we have

$$\mathbb{E}(|(f(X_0) - f(X'_0))(f(X_k) - f(X'_k))| | X_0, X'_0) \leq (c \cdot \ell_0)(c \cdot \ell_k).$$

Note that with $(X_0, X'_0) = (X, X')$, X_0 and X'_0 can differ at most at one coordinate. Suppose they differ at coordinate i . Then

$$(c \cdot \ell_0)(c \cdot \ell_k) \leq (c \cdot \ell_0)(c \cdot B^k \ell_0) = c_i(c \cdot B^k e_i),$$

where e_i denotes the vector whose i^{th} coordinate is 1 and all other coordinates are 0. Now, the probability that the X_0 and X'_0 differ at the i^{th} coordinate is $\leq 1/n$. Hence,

$$\begin{aligned} & \mathbb{E}(|(f(X_0) - f(X'_0))(f(X_k) - f(X'_k))| | X_0) \\ & \leq \frac{1}{n} \sum_{i=1}^n c_i (c \cdot B^k e_i) = \frac{1}{n} c \cdot B^k c \leq \frac{\|B\|_2^k \|c\|^2}{n} \end{aligned}$$

Thus, by (4.7) and (4.6), we get

$$v(X) \leq \frac{\|c\|^2}{2n(1 - \|B\|_2)} \leq \frac{\|c\|^2}{2(1 - \|A\|_2)}.$$

The proof is now completed by using Theorem 3.3. $\square$

4.3 Example: Spin systems

We now specialize to the case $\Omega = [-1, 1]$. Let ν be a probability measure on $[-1, 1]$. Suppose our original measure μ on $[-1, 1]^n$ has a density with respect to ν^n , represented in the Boltzmann form: $Z^{-1} e^{-H(x)}$, where Z is the normalizing constant.

The measure ν will usually be either the normalized Lebesgue measure, or the symmetric distribution on $\{-1, 1\}$. The generalized framework will allow us to deal with both discrete and continuous problems.

The following lemma gives us a simple way of applying Theorem 4.3 in this setting under the assumption that H can be extended to a twice continuously differentiable function on $[-1, 1]^n$.

Lemma 4.4 *For each pair (i, j) with $i \neq j$, define*

$$a_{ij} := 4 \sup_{x \in [-1, 1]^n} \left| \frac{\partial^2 H}{\partial x_i \partial x_j}(x) \right|$$

and let $a_{ii} = 0$ for each i . Then for each i and $x, y \in [-1, 1]^n$, we have

$$d_{TV}(\mu_i(\cdot | \bar{x}^i), \mu_i(\cdot | \bar{y}^i)) \leq \sum_{j=1}^n a_{ij} \mathbb{I}\{x_j \neq y_j\}.$$

Remark. It is obvious that this lemma, combined with Theorem 4.3, gives concentration inequalities for most of the familiar spin models at sufficiently high temperature. It covers, for instance, the Ising ferromagnets (in any dimension), the xy-model, and even the Ising spin glasses. This lemma is in fact, just a more easily recognizable version (for probabilists) of Proposition 8.8 in Georgii's book [45], which gives a similar condition for Gibbs potentials to satisfy Dobrushin's condition, and gives some more examples.

Example. Consider the Ising model on a graph $G = (V, E)$, where $V = \{1, \dots, n\}$ and the maximum degree is r . Here $\Omega = \{-1, 1\}$ and ν is the symmetric probability distribution on Ω . The Hamiltonian at inverse temperature β and external field h is given by

$$H_\beta(x) := -\beta \sum_{(i,j) \in E} x_i x_j - \beta h \sum_i x_i.$$

We shall consider the concentration of the *magnetization*, defined as

$$m(x) := \frac{1}{n} \sum_{i=1}^n x_i.$$

The Hamiltonian has a natural extension as a C^2 function on $\mathbb{R}^n$, and a simple computation gives

$$a_{ij} = \begin{cases} 4\beta & \text{if } (i, j) \in E, \\ 0 & \text{if } (i, j) \notin E. \end{cases}$$

Thus, the L^∞ and L^1 norms of the matrix $A = (a_{ij})$ are both bounded by $4\beta r$, where recall that r is the maximum degree of G . Thus, $\|A\|_2 \leq \sqrt{\|A\|_1 \|A\|_\infty} \leq 4\beta r$.

Again, it is clear that

$$|m(x) - m(y)| \leq \frac{2}{n} \sum_{i=1}^n \mathbb{I}\{x_i \neq y_i\}$$

Thus, if $X = (X_1, \dots, X_n)$ is a spin configuration drawn from the Gibbs measure at inverse temperature $\beta < 1/4r$ and external field h , and $m(X)$ is its magnetization,

then by Lemma 4.4 and Theorem 4.3, we have

$$\mathbb{P}\{|m(X) - \mathbb{E}m(X)| \geq t\} \leq e^{-\frac{1}{4}n(1-4\beta r)t^2}.$$

This gives an explicit concentration bound on the magnetization at sufficiently high temperature, though it need not necessarily cover the entire high temperature domain.

Proof of Lemma 4.4. The density with respect to ν of the conditional distribution $\mu_i(\cdot|\bar{x}^i)$ is given by

$$\rho_i(u|\bar{x}^i) = \frac{e^{-H(u, \bar{x}^i)}}{\int e^{-H(v, \bar{x}^i)} \nu(dv)},$$

where $(u, \bar{x}^i)$ denotes the vector obtained by substituting the number u as the i^{th} coordinate of the vector x .

Now fix i and j , where $i \neq j$. Direct computation shows that

$$\frac{\partial}{\partial x_j} \rho_i(u|\bar{x}^i) = - \left(\frac{\partial H}{\partial x_j}(u, \bar{x}^i) - \int \frac{\partial H}{\partial x_j}(v, \bar{x}^i) \rho_i(v|\bar{x}^i) \nu(dv) \right) \rho_i(u|\bar{x}^i)$$

and hence

$$\begin{aligned} \left| \frac{\partial}{\partial x_j} \rho_i(u|\bar{x}^i) \right| &\leq \sup_v \left| \frac{\partial H}{\partial x_j}(u, \bar{x}^i) - \frac{\partial H}{\partial x_j}(v, \bar{x}^i) \right| \rho_i(u|\bar{x}^i) \\ &\leq \frac{a_{ij}}{2} \rho_i(u|\bar{x}^i). \end{aligned}$$

Thus, for any $A \subseteq [-1, 1]$, we have

$$\left| \frac{\partial}{\partial x_j} \mu_i(A|\bar{x}^i) \right| = \left| \int_A \frac{\partial}{\partial x_j} \rho_i(u|\bar{x}^i) \nu(du) \right| \leq \frac{a_{ij}}{2}$$

Thus, we can conclude that for any $x, y \in [-1, 1]^n$,

$$\begin{aligned} d_{TV}(\mu_i(\cdot|\bar{x}^i), \mu_i(\cdot|\bar{y}^i)) \\ = \sup_A |\mu_i(A|\bar{x}^i) - \mu_i(A|\bar{y}^i)| \leq \sum_{j=1}^n a_{ij} \mathbb{I}\{x_j \neq y_j\}. \end{aligned}$$

This completes the argument. □

4.4 Example: Graph colorings

Suppose $G = (V, E)$ is a graph with n vertices and maximum degree r . A k -coloring of G is an assignment of k colors to the vertices of G . In other words, it is an element of $\{1, \dots, k\}^G$. A *proper* k -coloring is a k -coloring of G in which no two adjacent vertices receive the same color.

Let $X = (X_i, i \in V)$ be a coloring of G chosen uniformly from the set of all proper k -colorings.

A substantial amount of recent effort in theoretical computer science and associated probability theory has been devoted to the study of this random variable X and Markov chains converging in distribution to X . Most of it centers around temporal and spatial mixing and decay of correlations. An early observation of Jerrum [56] and Salas & Sokal [96], followed by vigorous activity in the last few years, have resulted in a spate of sophisticated coupling techniques and improved results. For up-to-date references, one can look at the recent articles [1, 39].

However, although coupling techniques have revolutionized the analysis of spatial and temporal decay of correlations in graph colorings, it is still extremely difficult to get concentration inequalities for general functions in these settings. In fact, we can see no direct way of getting concentration bounds for as simple a functional as the proportion of vertices having a particular color using currently available techniques.

In this section, we shall use our methods to get concentration inequalities for arbitrary (generalized) Lipschitz functions of randomly chosen proper graph colorings. That is, we shall consider $f : \{1, \dots, k\}^G \rightarrow \mathbb{R}$ satisfying $|f(x) - f(y)| \leq \sum_{i=1}^n c_i \mathbb{I}\{x_i \neq y_i\}$, and get tail bounds on $f(X)$, where X is our random proper coloring of G .

Both Jerrum [56] and Salas & Sokal [96] use the greedy coupling to get decay of correlations under the condition $k > 2r$. This is sort of a “high temperature phase” for proper graph colorings. Since our purpose is only to demonstrate the manner of application of our technique, we shall stick to this primitive approach.

Proposition 4.5 *Suppose G is a finite graph with maximum degree r , X is a uniformly chosen proper k -coloring of G , and $f : \{1, \dots, k\}^G \rightarrow \mathbb{R}$ satisfies $|f(x) - f(y)| \leq \sum_{i=1}^n c_i \mathbb{I}\{x_i \neq y_i\}$. If $k > 2r$, then for any $t \geq 0$ we have*

$$\mathbb{P}\{|f(X) - \mathbb{E}f(X)| \geq t\} \leq 2 \exp\left(-\frac{\gamma t^2}{\sum_{i=1}^n c_i^2}\right),$$

where $\gamma = (k - 2r)/(k - r)$.

Example 1. Let $f(x)$ be the proportion of vertices receiving color 1. It's clear that

$$|f(x) - f(y)| \leq \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{x_i \neq y_i\}.$$

Also, by symmetry, $\mathbb{E}f(X) = 1/k$, where X is drawn uniformly from the set of all proper k -colorings. Using the above result, we get

$$\mathbb{P}\{|f(X) - k^{-1}| \geq t\} \leq 2e^{-n\gamma t^2},$$

where $\gamma = (k - 2r)/(k - r)$ as in the lemma.

Example 2. Let $f(x)$ be the number of neighbors of vertex 1 which receive color 1. Then we can take $c_i = 1$ if i is a neighbor of 1 and $c_i = 0$ otherwise. Consequently, $\sum_i c_i^2 = r_1$, where r_1 is the degree of vertex 1. Thus,

$$\mathbb{P}\{|f(X) - \mathbb{E}f(X)| \geq t\} \leq 2e^{-\gamma t^2/r_1}.$$

Proof of Proposition 4.5. As usual, denote the law of X_i given $\bar{X}^i$ by $\mu_i(\cdot | \bar{x}^i)$. It is easy to see from definition that $\mu_i(\cdot | \bar{x}^i)$ is the uniform distribution over the set

$$\{1, \dots, k\} \setminus \{x_j : (i, j) \in E\}.$$

Thus, given two vectors x and y which differ only at a coordinate j , where j is adjacent

to i , it is clear that

$$\begin{aligned} d_{TV}(\mu_i(\cdot|\bar{x}^i), \mu_i(\cdot|\bar{y}^i)) &= \frac{1}{2} \sum_{u=1}^k |\mathbb{P}\{X_i = u|\bar{X}^i = \bar{x}^i\} - \mathbb{P}\{X_i = u|\bar{X}^i = \bar{y}^i\}| \\ &\leq \frac{1}{2} (\mathbb{P}\{X_i = y_j|\bar{X}^i = \bar{x}^i\} + \mathbb{P}\{X_i = x_j|\bar{X}^i = \bar{y}^i\}) \\ &\leq \frac{1}{k-r}. \end{aligned}$$

Thus, we can take $a_{ij} = 1/(k-r)$ if $(i, j) \in E$ and $a_{ij} = 0$ otherwise, and apply Theorem 4.3 to complete the proof. $\square$

4.5 Concentration of Haar measures

Let G be a compact topological group. Then there exists a G -valued random variable X with the properties that for any $x \in G$, the random variables xX , Xx and X^{-1} all have the same distribution as X . As mentioned in section 2.6, where we review the literature about concentration of measure on groups, the existence and uniqueness of the distribution of X (which is called the “Haar measure on G ”), is a classical result (see e.g. Rudin [93], Theorem 5.14). Our goal in this section is to study the concentration of the Haar measure using the convergence properties of certain kinds of Markov chains which have the Haar measure as their stationary distribution. An original application to the concentration of the spectra of sums of random matrices (related to free probability) will be given in the next section.

We shall not repeat the discussion of the existing literature on concentration of Haar measures, which is done in section 2.6 of the review chapter. It suffices to say that there is not much work. The only general theorem we could detect is a result of Milman & Schechtman [77] which is stated in section 2.6, where we also discuss available results for special groups like SO_n (the special orthogonal group) and S_n (the symmetric group).

We should, however, mention that we are not the first to analyze measures on groups using Stein’s method. Diaconis [27] has an application of Stein’s method to the analysis of random walks on groups. More recently, Jason Fulman has applied

Stein's method to analyze the Haar measure [43] and the Plancherel measure [44] on S_n .

Let us now introduce our setting. Let Y be a G -valued random variable having the following properties:

1. The random variable Y^{-1} has the same distribution as Y ; that is, the law of Y is *symmetric*.
2. For any $x \in G$, xyx^{-1} has the same distribution as Y . In other words, the distribution of Y is uniform on each conjugacy class of G .

Any such Y defines a reversible Markov kernel P in a natural way: For any $f : G \rightarrow \mathbb{R}$ such that $\mathbb{E}|f(X)| < \infty$, let

$$Pf(x) := \mathbb{E}f(Yx) = \mathbb{E}f(xx^{-1}Yx) = \mathbb{E}f(xY). \quad (4.8)$$

The reversibility of this kernel can be proved as follows: Since yX has the same distribution as X for any $y \in G$, therefore Y and YX are independent. Also, Y^{-1} has the same distribution as Y . Hence, the pair (X, Y) has the same distribution as (YX, Y^{-1}) . Consequently, the pairs (X, YX) and $(YX, Y^{-1}YX) = (YX, X)$ also have the same distribution. In other words, if we let $X' = YX$, the (X, X') is an exchangeable pair. Finally, note that $Pf(x) = \mathbb{E}(f(X')|X = x)$.

We shall attempt to study the concentration of X using the properties of this kernel P . A version of this problem was considered by Schmuckenschläger [98], but no practical solution was given. We shall now present a theorem that should suffice in many problems, but with “an extra $\log n$ factor”. As mentioned before, concrete examples will be provided later on.

Theorem 4.6 *Let G, X, Y be as above. Let $f : G \rightarrow \mathbb{R}$ be a measurable function such that $\mathbb{E}f(X) = 0$. Let $\|f\|_\infty = \sup_{x \in G} |f(x)|$ and*

$$\|f\|_Y := \sup_{x \in G} [\mathbb{E}(f(x) - f(Yx))^2]^{1/2}.$$

Let $Y_1, Y_2, \dots$, be i.i.d. copies of Y . Suppose a and b are two positive constants such that $d_{TV}(Y_1 Y_2 \cdots Y_k, X) \leq a e^{-bk}$ for every k , where d_{TV} is the total variation metric. Let A and B be two numbers such that $\|f\|_\infty \leq A$ and $\|f\|_Y \leq B$. Let

$$C = \frac{B^2}{2b} \left[\left(\log \frac{4aA}{B} \right)^+ + \frac{b(2 - e^{-b})}{1 - e^{-b}} \right].$$

Then $\text{Var}(f(X)) \leq C$, and for any $t \geq 0$, $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-t^2/2C}$.

Remarks. The rate of decay of the total variation distance for random walks on groups has been a topic of interest ever since the work of Diaconis & Shahshahani [33], who introduced group representation tools to overcome the deficiencies of the usual spectral gap approach (which gives a wrong answer for the random walk with transpositions on S_n). The property that Y is uniformly distributed on conjugacy classes is particularly suited to this line of analysis.

Since the seminal paper [33], much work has been done; several other methods have been developed by various authors, and the rates have been evaluated in many interesting problems. For a recent survey with an extensive bibliography, we refer to Diaconis [28].

To the best of our knowledge, Theorem 4.6 is the only result as of now, which connects the rates of convergence of random walks on groups — which is a widely explored area — with the concentration of Haar measures, which is not so widely explored.

Before we move on to complete the proof of Theorem 4.6, it is time for an easy application, which will not give a better-than-existing result, but merely demonstrate how Theorem 4.6 can be applied. An original application will be given in the next section.

Random permutations. Let $G = S_n$, the group of all permutations of n elements. Then the Haar measure is just the uniform distribution on G . Let X be a uniformly distributed random variables on G . We define the distribution of our Y by putting mass $1/n$ on the identity permutation and $2/n^2$ on each transposition of two elements. Since the set of transpositions is closed under conjugation and inversion, Y satisfies

the required properties 1 and 2 from the previous section. The kernel P defined by Y as in (4.8) is a familiar object which has been studied by several authors. The study was initiated by Diaconis & Shahshahani [33], who proved the following result:

Theorem 4.7 [Diaconis & Shahshahani [33]] *Let $Y_1, Y_2, \dots$, be i.i.d. copies of the random variable Y defined above. Then $d_{TV}(Y_1 \cdots Y_k, X) \leq 6ne^{-2k/n}$ for every $k > \frac{1}{2}n \log n$.*

The above version of the result has been quoted from [28], where it is written in a slightly different manner. Note that if we substitute $k = \frac{1}{2}n \log n$, the right hand side becomes 6, which is ≥ 1 , and hence the condition $k > \frac{1}{2}n \log n$ is redundant.

Using the Diaconis-Shahshahani result, we get $a = 6n$ and $b = 2/n$ in Theorem 4.6, and hence the following result:

Proposition 4.8 *Let $G = S_n$, where $n \geq 2$, and let X, Y be as above. Let $f : G \rightarrow \mathbb{R}$ be a function such that $\mathbb{E}f(X) = 0$. As in Theorem 4.6, let $\|f\|_Y = \max_{\sigma \in G} [\mathbb{E}(f(\sigma) - f(Y\sigma))^2]^{1/2}$. Let A and B be two numbers such that $\|f\|_\infty \leq A$ and $\|f\|_Y \leq B$. Let*

$$C = \frac{nB^2}{4} \left[\left(\log \frac{24nA}{B} \right)^+ + \frac{(2/n)(2 - e^{-2/n})}{1 - e^{-2/n}} \right].$$

Then $\text{Var}(f(X)) \leq C$, and for any $t \geq 0$, we have $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-t^2/2C}$.

For a simple application of this proposition, consider the *descent function* on S_n , defined as

$$D(\sigma) = \sum_{i=1}^{n-1} \mathbb{I}\{\sigma(i) > \sigma(i+1)\}.$$

The number of descents of a permutation is an interesting quantity from statistical and combinatorial points of view. They have been studied extensively by Foata and Schutzenberger [42], Knuth [59] and in the unpublished notes of Diaconis & Pitman [31]. Fulman [43] has applied Stein's method to the study of descents.

Let X be uniformly distributed on S_n . Clearly, $\|D(X) - \mathbb{E}D(X)\|_\infty \leq n$. Now, for any $x \in G$ and any transposition y , $|D(x) - D(yx)| \leq 4$. Putting $A = n$ and $B = 4$ in Theorem 4.8, and assuming $n \geq 10$, we get $C \leq 4n(2 \log n + 3.1)$. Thus, for

$n \geq 10$, we get $\text{Var}(D(X)) \leq 4n(2 \log n + 3.1)$ and for any $t \geq 0$,

$$\mathbb{P}\{|D(X) - \mathbb{E}D(X)| \geq t\} \leq 2 \exp\left(-\frac{t^2}{8n(2 \log n + 3.1)}\right).$$

The bounds are clearly off by a factor of $\log n$, but that will be a perpetual inconvenience of using Theorem 4.6. However, it is interesting to note that Maurey's theorem ([70], stated as Theorem 2.14 in section 2.6) gives the bound $\mathbb{P}\{|D(X) - \mathbb{E}D(X)| \geq t\} \leq 2e^{-t^2/256n}$ in this problem, which, though technically "better" than our bound, is always going to be worse in all practical situations.

That apart, there is a more serious reason why Proposition 4.8 may give better results in some problems. Maurey's result essentially uses $\|f\|_{\text{Lip}}$ instead of $\|f\|_Y$, where $\|f\|_{\text{Lip}} = \max\{|f(\sigma) - f(\tau\sigma)| : \sigma \in S_n, \tau \text{ is a transposition}\}$. Clearly, $\|f\|_Y \leq \|f\|_{\text{Lip}}$, and the difference may be significant in some situations.

Talagrand's result ([107], Theorem 5.1), on the other hand, will probably dominate Proposition 4.8 most of the time, though it is not clear whether Talagrand's method can be used to derive a result based on something like $\|f\|_Y$.

We now move on to prove the central result of this section.

Proof of Theorem 4.6. Let $X' = YX$. As observed before, (X, X') is an exchangeable pair. Recall that we defined $Pf(x) = \mathbb{E}f(Yx)$. By Lemma 4.1,

$$v(x) \leq \frac{1}{2} \sum_{k=0}^{\infty} \mathbb{E}|(f(x) - f(Yx))(P^k f(x) - P^k f(Yx))|, \quad (4.9)$$

where $v(x)$ is the usual quantity in our theory, as defined in (3.1). The criterion (4.3) required for Lemma 4.1 holds, since for any $z \in G$,

$$\begin{aligned} |P^k f(z)| &= |P^k f(z) - \mathbb{E}f(X)| = |\mathbb{E}f(Y_1 \cdots Y_k z) - \mathbb{E}f(Xz)| \\ &\leq 2\|f\|_{\infty} d_{TV}(Y_1 \cdots Y_k, X) \leq 2\|f\|_{\infty} a e^{-bk}. \end{aligned}$$

This observation also gives

$$\begin{aligned} & \mathbb{E}|(f(x) - f(Yx))(P^k f(x) - P^k f(Yx))| \\ & \leq 4\|f\|_\infty a e^{-bk} \mathbb{E}|f(x) - f(Yx)| \leq 4\|f\|_\infty a e^{-bk} \|f\|_Y. \end{aligned} \quad (4.10)$$

Now recall the assumption 2 that for any $y \in G$, $y^{-1}Yy$ has the same distribution as Y . Thus, for any $x, y \in G$,

$$Pf(yx) = \mathbb{E}(Yyx) = \mathbb{E}(yy^{-1}Yyx) = \mathbb{E}(yYx).$$

So, if we let Y' be an independent copy of Y , then

$$\begin{aligned} \mathbb{E}(Pf(x) - Pf(Yx))^2 &= \mathbb{E}(\mathbb{E}(f(Y'x) - f(YY'x)|Y)^2) \\ &\leq \mathbb{E}(f(Y'x) - f(YY'x))^2 \\ &\leq \sup_{y' \in G} \mathbb{E}(f(y'x) - f(Yy'x))^2 = \|f\|_Y^2. \end{aligned}$$

Thus, $\|Pf\|_Y \leq \|f\|_Y$. Continuing by induction, we get $\|P^k f\|_Y \leq \|f\|_Y$. Thus,

$$\begin{aligned} & \mathbb{E}|(f(x) - f(Yx))(P^k f(x) - P^k f(Yx))| \\ & \leq (\mathbb{E}(f(x) - f(Yx))^2)^{1/2} (\mathbb{E}(P^k f(x) - P^k f(Yx))^2)^{1/2} \\ & \leq \|f\|_Y \|P^k f\|_Y \leq \|f\|_Y^2. \end{aligned} \quad (4.11)$$

Using (4.10) and (4.11) in (4.9), we get

$$\begin{aligned} v(x) &\leq \frac{1}{2} \sum_{k=0}^{\infty} \min\{\|f\|_Y^2, 4a\|f\|_\infty \|f\|_Y e^{-bk}\} \\ &\leq \frac{1}{2} \sum_{k=0}^{\infty} \min\{B^2, 4aABe^{-bk}\} \\ &= \frac{B^2}{2} \sum_{k=0}^{\infty} \min\{1, 4aAB^{-1}e^{-bk}\}. \end{aligned} \quad (4.12)$$

We shall now compute a bound on the above sum. For ease of notation let $\beta =$

$4aAB^{-1}$, and let $\gamma = b^{-1} \log \beta$. If $\beta < 1$, the sum is just a geometric series which is easy to evaluate. Now assume $\beta \geq 1$. Then γ is nonnegative. Now, an easy verification shows that $\beta e^{-b\gamma} = 1$, and $1 \geq \beta e^{-bk}$ if and only if $k \geq \gamma$. Hence,

$$\begin{aligned} \sum_{k=0}^{\infty} \min\{1, \beta e^{-bk}\} &\leq \gamma + 1 + \sum_{k \geq \gamma} \beta e^{-bk} \\ &\leq \gamma + 1 + \beta e^{-b\gamma} \sum_{r=0}^{\infty} e^{-br} = \gamma + 1 + \frac{1}{1 - e^{-b}}. \end{aligned}$$

To finish, we substitute this bound in (4.12), and use Theorems 3.2 and 3.3 from Chapter 3. $\square$

4.6 Application to random matrices and free probability

Let M be an $n \times n$ complex hermitian (i.e. self-adjoint) matrix. We shall fix the following notation for the rest of this section.

- The *empirical spectral measure* M is the probability measure on $\mathbb{R}$, denoted by μ_M , which puts $1/n$ on each eigenvalue of M , repeated by multiplicities.
- The empirical distribution function of M , denoted by F_M , is the distribution function corresponding to the empirical spectral measure.
- Let Δ be a diagonal matrix whose diagonal elements are the eigenvalues of M . Then any hermitian matrix which has the same spectrum as M can be written as $U\Delta U^*$, where U is a unitary matrix. Thus, a uniform measure on the set of all such matrices is naturally induced by the Haar measure on the group of all unitary matrices of order n , which we shall denote by $\mathcal{U}_n$. This probability measure will be denoted by ρ_M .

A fundamental observation of Voiculescu [114] is the following: If M and N are two hermitian matrices of order n with empirical measures μ_M and μ_N , and n is large,

then the empirical measure μ_{M+N} of $M + N$ is “approximately determined” by μ_M and μ_N (irrespective of M and N) for “most choices of M and N ”. The meaning of the phrases in quotes is made precise by the following result:

Theorem 4.9 [Voiculescu’s result, as stated in Biane [14]] *For each positive integer n , let A_n and B_n be two hermitian matrices, whose eigenvalues are bounded uniformly in n . Let μ_1 and μ_2 be two probability measures with compact supports on $\mathbb{R}$, such that $\mu_{A_n} \rightarrow \mu_1$ and $\mu_{B_n} \rightarrow \mu_2$ weakly as $n \rightarrow \infty$. Then there exists a probability measure, depending only on μ_1 and μ_2 , denoted by $\mu_1 \boxplus \mu_2$, such that $\mu_{A'_n+B'_n} \rightarrow \mu_1 \boxplus \mu_2$, whenever A'_n and B'_n are random matrices chosen independently with distributions ρ_{A_n} and ρ_{B_n} .*

Voiculescu [114] proved this striking result in a limiting form, using the method of moments and some concentration results of Szarek [106] and Gromov & Milman [47]. The theorem was used by Voiculescu to establish a connection between free probability theory and random matrices, which resulted in an explosion of activity in the free probability literature. Another proof of Voiculescu’s observation, using Stieltjes transforms, was given by Pastur and Vasilchuk [83]. A good review of the literature, as well as a good exposition, is given in Biane [14]. Another useful reference is the lecture notes by Voiculescu [115].

To the best of our knowledge, all the existing results are limiting statements; quantitative bounds on the concentration of F_{M+N} given F_M and F_N for finite n are not available in the literature. We shall now state and prove one such result, as an application of the machinery developed in the previous section.

Theorem 4.10 *Let Δ_1 and Δ_2 be two $n \times n$ real diagonal matrices. Let U and V be independent Haar distributed random elements of $\mathcal{U}_n$, the group of all unitary matrices of order n . Let*

$$H = U\Delta_1U^* + V\Delta_2V^*,$$

and let F_H be the empirical distribution function of H . Then, for every $x \in \mathbb{R}$, $\text{Var}(F_H(x)) \leq \kappa n^{-1} \log n$. where κ is a universal constant not depending on n , Δ_1 ,

Δ_2 or x . Moreover, we also have the concentration inequality

$$\mathbb{P}\{|F_H(x) - \mathbb{E}(F_H(x))| \geq t\} \leq 2 \exp\left(-\frac{nt^2}{2\kappa \log n}\right)$$

for every $t \geq 0$, where κ is the same as in the variance bound.

To prove Theorem 4.10, we first need to establish a theorem about the concentration of the Haar measure on $\mathcal{U}_n$. Existing results of the type discussed in section 2.6 cannot give concentration bounds for F_H , since they are based on the Hilber-Schmidt distance which is too crude for such a delicate problem. Instead, we shall try to find the concentration of the Haar measure with respect to the *rank distance*, defined as $d(M, N) := \text{rank}(M - N)$. That this is indeed a metric, follows from the subadditivity of rank. The empirical distribution function is well-behaved with respect to this metric, as shown by the following lemma of Bai [8]:

Lemma 4.11 [Bai [8], Lemma 2.2] *Let M and N be two $n \times n$ hermitian matrices, with empirical distribution functions F_M and F_N . Then*

$$\|F_M - F_N\|_\infty \leq \frac{1}{n} \text{rank}(M - N).$$

This lemma is an easy consequence of the interlacing inequalities for eigenvalues of hermitian matrices. It seems possible that this already existed in the literature before Bai [8], but we could not find any reference.

We shall use Theorem 4.6 to find the concentration of the Haar measure on $\mathcal{U}_n$ with respect to the rank metric. For that purpose, we need a random walk which takes “small steps” with respect to that metric.

Let $G = \mathcal{U}_n$ and X be a Haar-distributed random variable on $\mathcal{U}_n$. We define the Y required for generating the random walk for Theorem 4.6 as follows: Let $Y = I - (1 - e^{i\varphi})uu^*$, where u is drawn uniformly from the unit sphere in $\mathbb{C}^n$, and φ is drawn independently from the distribution on $[0, 2\pi)$ with density proportional to $(\sin(\varphi/2))^{n-1}$. Multiplication by Y represents reflection across a randomly chosen

subspace. It is easy to verify that $Y \in \mathcal{U}_n$. Now, for any $U \in \mathcal{U}_n$,

$$UYU^* = I - (1 - e^{i\varphi})(Uu)(Uu)^*,$$

and Uu is again uniformly distributed over the unit sphere in $\mathbb{C}^n$. Also, $Y^{-1} = Y^* = I - (1 - e^{-i\varphi})uu^* = I - (1 - e^{i(2\pi-\varphi)})uu^*$ has the same distribution as Y , since $2\pi - \varphi$ has the same distribution as φ . Hence Y satisfies the properties 1 and 2 from section 4.5. Following a sketch of Diaconis & Shahshahani [34], Ursula Porod [88] proved the following result:

Theorem 4.12 [Porod [88]] *Let X, Y be as above. Let $Y_1, Y_2, \dots$, be i.i.d. copies of Y . There exists universal constants α, β, c_0 , such that whenever $n \geq 16$ and $k \geq \frac{1}{2}n \log n + c_0n$, we have*

$$d_{TV}(Y_1 \cdots Y_k, X) \leq \alpha n^{\beta/2} e^{-\beta k/n}. \quad (4.13)$$

Substituting $k = \frac{1}{2}n \log n + c_0n$, we get $\alpha e^{-\beta c_0}$ on the right hand side. Thus by suitably increasing α such that $\alpha e^{-\beta c_0} \geq 1$, we can drop the condition that $k \geq \frac{1}{2}n \log n + c_0n$. Combining Porod's theorem with Theorem 4.6, we get the following result about concentration of the Haar measure on $\mathcal{U}_n$:

Proposition 4.13 *Let $G = \mathcal{U}_n$ and X, Y be as above, with $n \geq 16$. Let $f : \mathcal{U}_n \rightarrow \mathbb{R}$ be a function such that $\mathbb{E}f(X) = 0$. Let $\|f\|_Y = \sup_{U \in \mathcal{U}_n} [\mathbb{E}(f(U) - f(YU))^2]^{1/2}$. Let A and B be constants such that $\|f\|_\infty \leq A$ and $\|f\|_Y \leq B$. Let*

$$C = \frac{nB^2}{2\beta} \left[\left(\log \frac{4\alpha n^\beta A}{B} \right)^+ + \frac{(\beta/n)(2 - e^{-\beta/n})}{1 - e^{-\beta/n}} \right],$$

where α and β are as in (4.13). Then $\text{Var}(f(X)) \leq C$, and for any $t \geq 0$, we have $\mathbb{P}\{|f(X)| \geq t\} \leq 2e^{-t^2/2C}$.

We are now ready to finish the proof of Theorem 4.10:

Proof of Theorem 4.10. We shall carry on with all the notation we have already defined in this section. The matrix $V^*HV = V^*U\Delta_1U^*V + \Delta_2$ has the same spectrum

as H . Also, V^*U is again Haar distributed. Hence, we can write, without loss of generality,

$$H = X\Delta_1X^* + \Delta_2,$$

where X follows the Haar distribution on $\mathcal{U}_n$. Now let

$$H' = (YX)\Delta_1(YX)^* + \Delta_2.$$

Recall that $Y = I - (1 - e^{i\varphi})uu^*$, where u is drawn from the uniform distribution on the unit sphere in $\mathbb{C}^n$, and φ is drawn independently from the distribution on $[0, 2\pi)$ with density proportional to $(\sin(\varphi/2))^{n-1}$. Let $\delta = 1 - e^{i\varphi}$. Then

$$\begin{aligned} H - H' &= X\Delta_1X^* - (I - \delta uu^*)X\Delta_1X^*(I - \bar{\delta}uu^*) \\ &= \delta H uu^* + \bar{\delta} uu^* H - |\delta|^2 uu^* H uu^*. \end{aligned}$$

The three summands are all of rank 1. It follows by the subadditivity of rank that $\text{rank}(H - H') \leq 3$. Thus by Lemma 4.11, we get

$$\|F_H - F_{H'}\|_\infty \leq \frac{3}{n}. \quad (4.14)$$

Now fix a point $x \in \mathbb{R}$, and let $f : \mathcal{U}_n \rightarrow \mathbb{R}$ be the map which takes X to $F_H(x)$. Then by (4.14), we have

$$|f(X) - f(YX)| \leq \frac{3}{n} \quad \text{for all possible values of } X \text{ and } Y.$$

Thus, $\|f\|_Y \leq 3/n$. Also, $\|f\|_\infty \leq 1$. Thus, in Proposition 4.13, we get $C \leq \kappa \log n + c$ for some universal constants κ and c . By choosing κ large enough, we can drop the assumption that $n \geq 16$ and also put $c = 0$. This completes the proof. $\square$

Bibliography

- [1] ACHLIOPTAS, D.; MOLLOY, M.; MOORE, C.; VAN BUSSEL, F. (2004). Sampling grid colorings with fewer colors. *Proc. 6th Latin American Theoretical Informatics (LATIN 2004)* 80-89.
- [2] ALON, N.; KRIVELEVICH, M.; VU, V. H. (2002). On the concentration of eigenvalues of random symmetric matrices. *Israel J. Math.* **131** 259–267.
- [3] AMIR, D.; MILMAN, V. D. (1980). Unconditional and symmetric sets in n -dimensional normed spaces. *Israel J. Math.* **37** No. 1-2, 3–20.
- [4] ANÉ, C.; BLACHÈRE, S.; CHAFAÏ, D.; FOUGÈRES, P.; GENTIL, I.; MALRIEU, F.; ROBERTO, C.; SCHEFFER, G. (2000). *Sur les inégalités de Sobolev logarithmiques*. Société Mathématique de France, Paris.
- [5] ARRATIA, R.; GOLDSTEIN, L.; GORDON, L. (1989). Two moments suffice for Poisson approximations: the Chen-Stein method. *Ann. Probab.* **17** No. 1, 9–25.
- [6] ARRATIA, R.; GOLDSTEIN, L.; GORDON, L. (1992). Poisson approximation and the Chen-Stein method. *Statist. Sci.* **5** No. 4, 403–434.
- [7] AZUMA, K. (1967). Weighted sums of certain dependent random variables. *Tôhoku Math. Journal* **19** 357–367
- [8] BAI, Z. D. (1999). Methodologies in spectral analysis of large-dimensional random matrices, a review. *Statist. Sinica* **9** No. 3, 611–677.

- [9] BAKRY, D.; ÉMERY, M. (1985). Diffusions hypercontractives. *Séminaire de probabilités, XIX, 1983/84*, Lecture Notes in Math., **1123**, Springer, Berlin, 177–206.
- [10] BARBOUR, A. D.; HOLST, L.; JANSON, S. (1992). *Poisson approximation*. Oxford Studies in Probability, **2** The Clarendon Press, Oxford University Press, New York.
- [11] BENNETT, G. (1962). Probability inequalities for sums of independent random variables. *J. Amer. Statist. Assoc.* **57** 33–45.
- [12] BESAG, J. E. (1972). Nearest-neighbour systems and the auto-logistic model for binary data. *J. Roy. Statist. Soc. Ser. B* **34** 75–83.
- [13] BESAG, J. E. (1975). Statistical analysis of non-lattice data. *The Statistician*, **24** 179–195.
- [14] BIANE, P. (2003). Free probability for probabilists. *Quantum probab. commun., Vol. XI (Grenoble, 1998)*, 55–71, QP-PQ, XI, World Sci. Publishing, River Edge, NJ.
- [15] BOBKOV, S.; GENTIL, I.; LEDOUX, M. (2001). Hypercontractivity of Hamilton–Jacobi equations. *J. Math. Pures Appl. (9)* **80** 669–696.
- [16] BOBKOV, S.; LEDOUX, M. (1997). Poincaré’s inequalities and Talagrand’s concentration phenomenon for the exponential distribution. *Probab. Theory Related Fields* **107** No. 3, 383–400.
- [17] BOBKOV, S.; GÖTZE, F. (1999). Exponential integrability and transportation cost related to logarithmic Sobolev inequalities. *J. Funct. Anal.* **163** 1–28.
- [18] BOUCHERON, S.; LUGOSI, G.; MASSART, P. (2000). A sharp concentration inequality with applications. *Random Structures Algorithms*. **16** 277–292.
- [19] BOUCHERON, S.; LUGOSI, G.; MASSART, P. (2003). Concentration inequalities using the entropy method. *Ann. Probab.* **31** No. 3, 1583–1614.

- [20] BOUCHERON, S.; BOUSQUET, O.; LUGOSI, G; MASSART, P. (2005). Moment inequalities for functions of independent random variables. *Ann. Probab.* **33** No. 2, 514–560.
- [21] BURKHOLDER, D. L. (1973). Distribution function inequalities for martingales. *Ann. Probab.* **1** 19–42.
- [22] BURKHOLDER, D. L. (1988). Sharp inequalities for martingales and stochastic integrals. *Astérisque*, **157-158** 75–94.
- [23] BURKHOLDER, D. L. (1989). Explorations in martingale theory and its applications. *Ecole d’Eté de Probabilités de Saint-Flour XIX*, 1–66. Springer, Lecture Notes in Mathematics #**1464**.
- [24] CHATTERJEE, S.; DIACONIS, P.; MECKES, E. (2005). Exchangeable pairs and Poisson approximation. *Probab. Surv.* **2** 64–106.
- [25] CHEN, L. H. Y. (1975). Poisson approximation for dependent trials. *Ann. Probab.* **3** No. 3, 534–545.
- [26] DEMBO, A. (1997). Information inequalities and concentration of measure. *Ann. Probab.* **25** 927–939.
- [27] DIACONIS, P. (1991). An example for Stein’s method. *Stanford Univ. Tech. Report.* #256384.
- [28] DIACONIS, P. (2003). Random walks on groups: characters and geometry. *Groups St. Andrews 2001 in Oxford. Vol. I*, 120–142, London Math. Soc. Lecture Note Ser., **304**, Cambridge Univ. Press, Cambridge.
- [29] DIACONIS, P.; GRAHAM, R. L. (1977). Spearman’s footrule as a measure of disarray. *J. Roy. Statist. Soc. Ser. B* **39** No. 2, 262–268.
- [30] DIACONIS, P.; HOLMES, S. (EDITORS) (2004). *Stein’s method: expository lectures and applications*. IMS Lecture Notes—Monograph Series, **46**.
- [31] DIACONIS, P.; PITMAN, J. (1991). Unpublished notes on descents.

- [32] DIACONIS, P.; SALOFF-COSTE, L. (1996). Logarithmic Sobolev inequalities for finite Markov chains. *Ann. Appl. Probab.* **6** No. 3, 695–750.
- [33] DIACONIS, P.; SHAHSHAHANI, M. (1981). Generating a random permutation with random transpositions. *Z. Wahrsch. Verw. Gebiete* **57** No. 2, 159–179.
- [34] DIACONIS, P.; SHAHSHAHANI, M. (1986). Products of random matrices as they arise in the study of random walks on groups. *Random matrices and their applications (Brunswick, Maine, 1984)*, 183–195, Contemp. Math., **50**, Amer. Math. Soc., Providence, RI.
- [35] DOBRUSHIN, R. L. (1970). Prescribing a system of random variables by conditional distributions. *Theory Probab. Appl.* **15** 453–486.
- [36] DOBRUSHIN, R. L.; SHLOSMAN, S. (1985). Constructive criterion for the uniqueness of Gibbs field. In *Statistical Physics and Dynamical Systems* (J. Fritz, A. Jaffe, and D. Szász, eds.) 347–369. Birkhäuser, Basel.
- [37] DOBRUSHIN, R. L.; SHLOSMAN, S. (1987). Completely analytical interactions: Constructive description. *J. Statist. Phys.* **46** 983–1014.
- [38] DVORETZKY, A. (1961). Some results on convex bodies and Banach spaces. *Proc. Internat. Sympos. Linear Spaces, Jerusalem*. 123–160 Jerusalem Academic Press, Jerusalem; Pergamon, Oxford.
- [39] DYER, M.; FRIEZE, A.; HAYES, T. P.; VIGODA, E. (2004). Randomly coloring constant degree graphs. *Proc. 45th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2004)*.
- [40] EFRON, B.; STEIN, C. (1981). The jackknife estimate of variance. *Ann. Statist.* **9** No. 3, 586–596.
- [41] ELLIS, R. S. (1985). *Entropy, large deviations, and statistical mechanics*. Grund. der Mathemat. Wissenschaften, **271**. Springer-Verlag, New York.

- [42] FOATA, D.; SCHUTZENBERGER, M. P. (1970). Theorie geometriques des polynomes Euleriens. *Springer Lecture Notes in Math.* **138**.
- [43] FULMAN, J. (2004). Stein's method and non-reversible Markov chains. *Stein's method: expository lectures and applications*, 69–77, IMS Lecture Notes Monogr. Ser., **46**, IMS, Beachwood, OH.
- [44] FULMAN, J. (2005). Stein's method and Plancherel measure of the symmetric group. *Trans. Amer. Math. Soc.* **357** No. 2, 555–570 (electronic).
- [45] GEORGII, H.-O. (1988). *Gibbs measures and phase transitions*. de Gruyter Studies in Mathematics, **9**. Walter de Gruyter & Co., Berlin.
- [46] GEYER, C. J.; THOMPSON, E. A. (1992). Constrained Monte Carlo maximum likelihood for dependent data. *J. Roy. Statist. Soc. Ser. B* **54** No. 3, 657–699.
- [47] GROMOV, M.; MILMAN, V. D. (1983). A topological application of the isoperimetric inequality. *Amer. J. Math.* **105** No. 4, 843–854.
- [48] GROSS L. (1976). Logarithmic Sobolev inequalities. *Amer. J. Math.* **97** 1061–1083.
- [49] GUIONNET, A.; ZEGARLINSKI, B. (2003). Lectures on logarithmic Sobolev inequalities. *Séminaire de Probabilités, XXXVI*, Lecture Notes in Math., **1801**, Springer, Berlin, 1–134.
- [50] GUIONNET, A.; ZEITOUNI, O. (2000). Concentration of the spectral measure for large matrices. *Electron. Comm. Probab.* **5** 119–136.
- [51] Hoeffding, W. (1951). A combinatorial central limit theorem. *Ann. Math. Statist.* **22** No. 4, 558–566.
- [52] Hoeffding, W. (1963). Probability inequalities for sums of bounded random variables. *J. Amer. Stat. Assoc.* **58** 13–30.

- [53] HOLLEY, R.; STROOCK, D. (1987). Logarithmic Sobolev Inequalities and Stochastic Ising Models. *J. Stat. Phys.* **46** No. 5-6, 1159–1194.
- [54] HOUDRÉ, C.; TETALI, P. (2001). Concentration of measure for products of Markov kernels and graph products via functional inequalities. *Combin. Probab. Comput.* **10** No. 1, 1–28.
- [55] HOUDRÉ, C.; TETALI, P. (2004). Isoperimetric invariants for product Markov chains and graph products. *Combinatorica* **24** No. 3, 359–388.
- [56] JERRUM, M. R. (1995). A very simple algorithm for estimating the number of k -colorings of a low-degree graph. *Random Struct. Algorithms*, **7** No. 2, 157–165.
- [57] KENDALL, M.; GIBBONS, J. D. (1990). *Rank correlation methods*. Fifth ed. Edward Arnold, London.
- [58] KIM, J. H.; VU, V. H. (2004). Divide and conquer martingales and the number of triangles in a random graph. *Random Struct. Algorithms* **24** No. 2, 166–174.
- [59] KNUTH, D. (1973). *The art of computer programming. Volume 3. Sorting and searching*. Addison-Wesley Publishing Co.
- [60] LEDOUX, M. Concentration of measure and logarithmic Sobolev inequalities. *Séminaire de Probabilités, XXXIII*, Lecture Notes in Math., **1709**, Springer, Berlin. 120–216.
- [61] LEDOUX, M. (2001). *The Concentration of Measure Phenomenon*. Amer. Math. Soc., Providence, RI.
- [62] LEDOUX, M. (2001). Logarithmic Sobolev inequalities for unbounded spin systems revisited. *Séminaire de Probabilités, XXXV*, 167–194, Lecture Notes in Math., **1755**, Springer, Berlin, 2001.
- [63] LÉVY, P. (1951). *Problèmes concrets d'analyse fonctionnelle*. Gauthier-Villars, Paris.

- [64] MAKOWSKI, A. M.; SHWARTZ, A. (2002). The Poisson equation for countable Markov chains: probabilistic methods and interpretations. *Handbook of Markov decision processes*, 269–303, Internat. Ser. Oper. Res. Management Sci., **40**, Kluwer Acad. Publ., Boston, MA.
- [65] MARTON, K. (1996). Bounding $\bar{d}$ -distance by information divergence: A method to prove measure concentration. *Ann. Probab.* **24** 857–866.
- [66] MARTON, K. (1996). A measure concentration inequality for contracting Markov chains. *Geom. Funct. Anal.* **6** 556–571.
- [67] MARTON, K. (2003). Measure concentration and strong mixing. *Studia Sci. Math. Hungar.* **40** No. 1–2, 95–113.
- [68] MARTON, K. (2004). Measure concentration for Euclidean distance in the case of dependent random variables. *Ann. Probab.* **32** 3B, 2526–2544.
- [69] MASSART, P. (2000). About the constants in Talagrand’s concentration inequalities for empirical processes. *Ann. Probab.* **28** 863–884.
- [70] MAUREY, B. (1979). Construction de suites symétriques. *C. R. Acad. Sci. Paris Sér. A-B* **288** No. 14, A679–A681.
- [71] MCDIARMID, C. (1989). On the method of bounded differences. *Surveys in combinatorics* (J. Siemons, ed.). *London Math. Soc. Lecture Notes Ser.* **141** 148–188.
- [72] MCDIARMID, C. (2002). Concentration for independent permutations. *Combin. Probab. Comput.* **11** No. 2, 163–178.
- [73] MEYER, P. A. (1971). Solutions de l’équation de Poisson dans le cas récurrent. *Séminaire de Probabilités, V*, 251–269. *Lecture Notes in Math.*, Vol. **191**, Springer, Berlin.
- [74] MÉZARD, M.; PARISI, G.; VIRASORO, M. A. *Spin glass theory and beyond*. World Scientific Publishing Co., Inc., Teaneck, NJ.

- [75] MILMAN, V. D. (1971). A new proof of A. Dvoretzky's theorem on cross-sections of convex bodies. *Funkcional. Anal. i Priložen.* **5** No. 4, 28–37.
- [76] MILMAN, V. D. (1988). The heritage of P. Lévy in geometrical functional analysis. Colloque Paul Lévy sur les Processus Stochastiques. *Astérisque* No. **157-158**, 273–301.
- [77] MILMAN, V. D.; SCHECHTMAN, G. (1986). Asymptotic theory of finite-dimensional normed spaces. Lecture Notes in Math., **1200**. Springer-Verlag, Berlin.
- [78] NEVEU, J. (1972). Potentiel Markovien récurrent des chaînes de Harris. *Ann. Inst. Fourier (Grenoble)* **22** 7–130.
- [79] NUMMELIN, E. (1991). On the Poisson equation in the potential theory of a single kernel. *Math. Scand.* **68** 59–82.
- [80] OTTO, F.; VILLANI, C. (2000). Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality. *J. Funct. Anal.* **173** 361–400.
- [81] OPPER, M.; SAAD, D. (EDITORS) (2001). *Advanced mean field methods. Theory and practice*. Neural Information Processing Series. MIT Press, Cambridge, MA.
- [82] PARISI, G. (1988). *Statistical field theory*. Benjamin/Cummings Publishing Co., Inc.
- [83] PASTUR, L.; VASILCHUK, V. (2000). On the law of addition of random matrices. *Comm. Math. Phys.* **214** No. 2, 249–286.
- [84] PESTOV, V. G. (2000). Amenable representations and dynamics of the unit sphere in an infinite-dimensional Hilbert space. *Geom. Funct. Anal.* **10** No. 5, 1171–1201.
- [85] PESTOV, V. G. (2000). Ramsey-Milman phenomenon, Urysohn metric spaces, and extremely amenable groups. *Israel J. Math.* **127** 317–357.

- [86] PICKARD, D. K. (1987). Inference for discrete Markov fields: the simplest nontrivial case. *J. Amer. Statist. Assoc.* **82** No. 397, 90–96.
- [87] POLLARD, D. (1990). *Empirical processes: theory and applications*. NSF-CBMS Regional Conference Ser. in Probab. and Statist., 2. IMS, Hayward, CA; ASA, Alexandria, VA.
- [88] POROD, U. (1996). The cut-off phenomenon for random reflections. II. Complex and quaternionic cases. *Probab. Theory Related Fields* **104** No. 2, 181–209.
- [89] REINERT, G. (1995). A weak law of large numbers for empirical measures via Stein’s method. *Ann. Probab.* **23** No. 1, 334–354.
- [90] REITZNER, M. (2003). Random polytopes and the Efron-Stein jackknife inequality. *Ann. Probab.* **31** No. 4, 2136–2166.
- [91] REVUZ, D. (1984). *Markov Chains*. 2nd ed. North-Holland, Amsterdam.
- [92] RIO, E. (2001). Inégalités de concentration pour les processus empiriques de classes de parties. *Probab. Theory Relat. Fields* **119** 163–175.
- [93] RUDIN, W. (1973). *Functional analysis*. McGraw-Hill Book Co., New York-Düsseldorf-Johannesburg.
- [94] RUDIN, W. (1976). *Principles of mathematical analysis*. Third edition. McGraw-Hill Book Co., New York-Auckland-Düsseldorf.
- [95] SAMSON, P.-M. (2000). Concentration of measure inequalities for Markov chains and Φ -mixing processes. *Ann. Probab.* **28** No. 1, 416–461.
- [96] SALAS, J.; SOKAL, A. D. (1997). Absence of phase transition for antiferromagnetic Potts models via Dobrushin’s uniqueness theorem. *J. Statistical Physics* **86** 551–579.
- [97] SCHECHTMAN, G. (1982). Lévy type inequality for a class of finite metric spaces. *Martingale theory in harmonic analysis and Banach spaces (Cleveland 1981)*. Lecture Notes in Math., **939** Springer. 211–215.

- [98] SCHMUCKENSCHLÄGER, M. (1998). Curvature of nonlocal Markov generators. In *Convex Geometric Analysis: MSRI Publications*. **34** 189–197.
- [99] SHAMIR, E.; SPENCER, J. (1987). Sharp concentration of the chromatic number on random graphs $G_{n,p}$. *Combinatorica* **7** No. 1, 121–129.
- [100] STEELE, J. M. (1986). An Efron-Stein inequality for nonsymmetric statistics. *Ann. Statist.* **14** 753–758.
- [101] STEIN, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. *Proc. of the Sixth Berkeley Symp. on Math. Statist. and Probab., Vol. II: Probability theory*. 583–602.
- [102] STEIN, C. (1986). *Approximate computation of expectations*. IMS Lecture Notes—Monograph Series, **7**.
- [103] STROOCK, D. W.; ZEGARLINSKI, B. (1992). The logarithmic Sobolev inequality for continuous spin systems on a lattice. *J. Funct. Anal.* **104** 299–326.
- [104] STROOCK, D. W.; ZEGARLINSKI, B. (1992). Equivalence of the logarithmic Sobolev inequality and the Dobrushin-Shlosman mixing condition. *Commun. Math. Phys.* **144** 303–323.
- [105] STROOCK, D. W.; ZEGARLINSKI, B. (1992). The logarithmic Sobolev inequality for discrete spin systems on a lattice. *Commun. Math. Phys.* **149** 175–193.
- [106] SZAREK, S. J. (1990). Spaces with large distance to l_∞^n and random matrices. *Amer. J. Math.* **112** No. 6, 899–942.
- [107] TALAGRAND, M. (1995). Concentration of measure and isoperimetric inequalities in product spaces. *Inst. Hautes tudes Sci. Publ. Math.* **81** 73–205.
- [108] TALAGRAND, M. (1996). Transportation cost for gaussian and other product measures. *Geom. Funct. Anal.* **6** 587–600.

- [109] TALAGRAND, M. (1996). New concentration inequalities in product spaces. *Invent. Math.* **126** 505–563.
- [110] TALAGRAND, M. (1996). A new look at independence. *Ann. Probab.* **24** 1–34.
- [111] TALAGRAND, M. (2003). *Spin glasses: a challenge for mathematicians. Cavity and mean field models*. Springer-Verlag, Berlin.
- [112] TALAGRAND, M. (2005). The Parisi formula. *Ann. Math.* To appear.
- [113] VILLANI, C. (2003). *Topics in optimal transportation*. Graduate Studies in Mathematics, 58. Amer. Math. Society, Providence, RI.
- [114] VOICULESCU, D. (1991). Limit laws for random matrices and free products. *Inv. Math.* **104** 201–220.
- [115] VOICULESCU, D. (2000). Lectures on free probability theory. *Lectures on probability theory and statistics (Saint-Flour, 1998)*, 279–349, Lecture Notes in Math., **1738**, Springer, Berlin.